JUL 24, 2020

SOLUTION- 12 – HBSS-HEPES for ROS measurement in PMN

Marco Cosentino¹, Elisa Storelli¹, Alessandra Luini¹,
Massimiliano LM Legnaro¹, Emanuela Rasini¹, Marco
Franca Marino¹, Ferrari¹,

¹Center for Research in Medical Pharmacology, University of Insubria
(Varese, Italy)

OPEN ACCESS

Farmacologia Medica

DOI:
dx.doi.org/10.17504/protocols.io.biyxkfxn

Document Citation: Marco Cosentino, Elisa Storelli, Alessandra Luini, Massimiliano LM Legnaro, Emanuela Rasini, Marco Ferrari, Franca Marino 2020. SOLUTION- 12 – HBSS-HEPES for ROS measurement in PMN. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.biyxkfxn>

License: This is an open access document distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Created: Jul 24, 2020

Last Modified: Jul 24, 2020

DOCUMENT integer ID:
39671

HBSS/HEPES x ROS composition (g/l):

NaCl: 4.2300 g

KCl: 0.1864 g

MgSO₄: 0.1232 g

CaCl₂: 0.735 g

Glucose: 0.900 g

Hepes: 1.300 g

To prepare 0.5 L of HBSS/HEPES, dissolve the reagents listed above in 450 mL of ultrapure H₂O.

Adjust the pH to 7.4 with HCl or NaOH, and then add H₂O to 0.5 L.

Storage: 4 °C Fridge 1- (Room TS08)

NaCl code: S9625, Sigma

KCl code: P9541, Sigma

MgSO₄ code: 1.05886, Sigma

CaCl₂ code: 1.02382 Sigma

Glucose code: 1.08337, Sigma

Hepes code: H7006 Sigma