MAY 20, 2020

Sorbitol washing complex homogenate for improved DNA extractions

PLOS One ✓ Peer-reviewed method

In 1 collection

Ashley Jones¹, Benjamin Schwessinger¹

¹Research School of Biology, Australian National University, Canberra, ACT, Australia

High molecular weight DNA extraction from all kingdoms

PLOS ONE Lab Protocols

OPEN ACCESS

DOI:

dx.doi.org/10.17504/protocols.io.beuvjew6

External link:

<https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0253830>

Protocol Citation: Ashley Jones, Benjamin Schwessinger 2020. Sorbitol washing complex homogenate for improved DNA extractions.

protocols.io

<https://dx.doi.org/10.17504/protocols.io.beuvjew6>

MANUSCRIPT CITATION:

Jones A, Torkel C, Stanley D, Nasim J, Borevitz J, Schwessinger B (2021) High-molecular weight DNA extraction, clean-up and size selection for long-read sequencing. PLoS ONE 16(7): e0253830.

doi.org/10.1371/journal.pone.0253830

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Ashley Jones

Australian National University

ABSTRACT

The extraction of pure DNA can be challenging due to the presence of sugars, oils and other endogenous chemicals present within biological samples. This is particularly true for many plants and fungi due to the presence of secondary metabolites such as polyphenols and polysaccharides. Polyphenols within the cytosol can become irreversibly DNA-bound after cell lysis and polysaccharides can co-precipitate with DNA during the extraction. Sorbitol is an osmotically active sugar alcohol and washing homogenate with sorbitol before cell lysis has been shown to significantly improve the purity of DNA extractions. Sorbitol does not pass cell membranes and likely acts by drawing the cytosol out of the cell. Therefore polyphenols and polysaccharides would be removed.

GUIDELINES

This protocol is based on the following publication. When citing, please also note the original publication below.

Inglis, P., Pappas, M.d.C., Resende, L. and Grattapaglia, D. (2018). Fast and inexpensive protocols for consistent extraction of high quality DNA and RNA from challenging plant and fungal samples for high-throughput SNP genotyping and sequencing applications. PLOS ONE 13, 1-14.

MATERIALS

Dithiothreitol (DTT) or β -Mercaptoethanol

EDTA pH 8

Milli-Q water (MQW)

PVP 40,000

D-Sorbitol

Tris-HCl pH 8

Created: Apr 09, 2020

Last Modified: May 20, 2020

PROTOCOL integer ID:
35445

PREPARATION

- 1 Prepare sorbitol wash solution. Approximately 2x volume of tube capacity will be used per sample. Sample tissue should not exceed 33% of tube capacity.

To prepare a 500 mL solution:

Reagent	Target concentration	Molecular weight	Stock concentration	Add from stock
D-Sorbitol	0.35 M	182.17	powder	31.88 g
PVP 40,000	1% (w/v)	40,000	powder	5 g
Tris-HCl pH 8	100 mM	157.60	1 M	50 mL
EDTA pH 8	5 mM	292.24	0.5 M	5 mL
MQW	NA	NA	NA	Bring to 500 mL

Note

- Store at 4°C for up to 6 months.
- DTT or β -Mercaptoethanol will be added freshly when used (below).

HOMOGENISATION

- 2 Grind tissue to a fine powder, keep frozen with liquid nitrogen.

- 3 If homogenate is not in a tube (i.e. mortar and pestle), transfer to an appropriate sized tube.

Note

Homogenate should not exceed 33% of tube capacity.

SORBITOL WASH

- 4 Fill the tube capacity with an excess of sorbitol wash solution.
- 5 Add DTT, approximate final concentration of 1 mM. Add 0.15 mg DTT per 1 mL of sorbitol wash. A tiny scoop of powder is sufficient. DTT MW = 154.253.

Note

Alternatively add 1% β -Mercaptoethanol (v/v). Add 10 μ L of β -Mercaptoethanol per 1 mL of sorbitol wash.

- 6 Shake, invert and vortex to mix thoroughly.

Note

Ensure tissue is in suspension.

- 7 Centrifuge at 5,000 rcf for 5 min at room temperature.

Note

Limited based on tubes rupturing if they contain ball bearings. Can also be reduced to 2,500 rcf.

- 8 Carefully decant the supernatant. Remove as much of the wash solution as possible without losing the pellet.

Note

Supernatant should appear slightly cloudy or discoloured light yellow or brown.

- 9 If the supernatant was turbid, viscous or had dark discolouration, repeat the sorbitol wash.

DNA EXTRACTION

- 10 Proceed to a DNA extraction method of choice by adding lysis buffer to the pellet. Continue protocol as normal.