MAY 18, 2021

SPARC Cat - Sham Control Chronic Cat 1 Day 30

Margot

Brett Hanzlicek¹, Anna Rietsch², Damaser²

¹Advanced Platform Technology Center, Louis Stokes VA Hospital, Cleveland, Ohio;

²Department of Biomedical Engineering, Cleveland Clinic Lerner Research Institute, Cleveland, Ohio

SPARC

Tech. support email: info@neuinfo.org

Brett Hanzlicek

ABSTRACT

This is a procedure for a sham control chronic cat experiment (Day 30-terminal) for cystotomy (bladder surgery). The cystotomy is performed without UroMOCA implantation. The cat is observed daily and imaged on day 0, 14 and 30 to track changes to the bladder and overall cat health in response to the cystotomy. This protocol includes urodynamics and imaging for Day 30 in the chronic experiments.

MATERIALS

Cat - Domestic short-haired cat. Male or female. 6-24 months; 3.5-5.0kg

Syringe Pump - Genie Touch; Kent Scientific

Pressure Transducer - Catalog # 503067

WPI Amplifier - 4 channel transbridge

WPI National Instruments multifunction I/O device - NI USB 6259

National Instruments Laptop with Labview software

Tubing - Masterflex L/S Platinum-Cured Silicone Tubing ID 3/16 "; Cole Parmer

BNC cables

Tubing to catheter - APIS

Laborie Catheter - Argyl Suction Catheter, 3.5 Fr Catalog # 8890703211

Covidien Sutures - 4-0 Vicryl Polysorb (Covidien UL212)

3-0 Prolene (Ethicon 8762)

2-0 Silk Syringe

60ml Contrast

Visipaque 320mg/ml

Surgilube

Surgical instruments

OPEN ACCESS

DOI:

dx.doi.org/10.17504/protocols.io.bfztjp6n

Protocol Citation: Brett Hanzlicek, Anna Rietsch, Margot Damaser 2021. SPARC Cat - Sham Control Chronic Cat 1 Day 30.

protocols.io

<https://dx.doi.org/10.17504/protocols.io.bfztjp6n>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: May 05, 2020

Last Modified: May 18, 2021

PROTOCOL integer ID:
36627

BEFORE START INSTRUCTIONS

1 week before start of experiment - Confirm absence of bladder spasm; healthy cat

12 hours before experiment, start fasting the cat

You must transport the animal chart along with the cat to surgery site. The veterinary team will record all relevant data in the chart. The chart must then go back to the housing site.

Transport Cat

- 1 Transport cat from housing site to surgery site.

Animal Prep and catheter placement

- 2 Animal is anesthetized and abdomen is shaved by the vet team. The cat is then moved into the surgery room and attached to monitors by the vet team.
- 3 Drape animal and perform betadine scrub on abdomen and genitals.
- 4 Put surgilube on 3.5Fr catheter and insert into bladder through the urethra. Advance the catheter until resistance is met, then pull back 2-3cm.
- 5 Use gentle suction with syringe to withdraw urine from bladder through the catheter. Measure the volume and save the urine for urinalysis.

15 mL urine removed

Cystogram - DYNA CT

- 6 Use 1:5 dilution of contrast to saline to visualize bladder.

Fill the bladder with 10 mL contrast and take a single CT image.

Add 10 mL more to bladder (total of 20ml) and take single CT image

Add 10 mL more to bladder (total of 30ml) and take single CT image

Take a 3D CT image with 30ml contrast:saline in the bladder

7 Empty Bladder

Urodynamics/Cystometry 1

8

Note

Performed after cat is transferred from Isoflurane to propofol anesthetic.

Note

Fill rate is 2ml saline/minute using syringe pump.

Note

Data is recorded using Labview software.

Note

Pressure is recorded using an external pressure transducer connected to syringe pump on one end and tubing that leads to the catheter on the other end.

9 Fill bladder with saline and record pressure using pressure transducer connected to LabView

Fill to 13.9 mL ; Peak pressure around 20 cmH₂O

Fill to 20 mL ; Peak pressure around 34 cmH₂O

Tug on catheter after filling to visualize contraction in Labview software

10 Empty Bladder

Urodynamics/cystometry 2

11 Fill bladder with saline at 2ml/min

12 Fill to 15 mL ; End round 2 urodynamics

Note

Appears bladder capacity is around 15-20 ml

Euthanasia

13 Cleveland Clinic animal team euthanizes the animal

Dissection/Histology

14 Bladder is removed, examined and placed in formalin for histological analysis