VERSION 3

JUN 30, 2020

OPEN ACCESS

DOI:

dx.doi.org/10.17504/protocols.io.bh37j8rn

Protocol Citation: Nicole A. Pelot, J. Ashley Ezzell, Gabriel B. Goldhagen, Jake E. Cariello, Kara A. Clissold, Warren M. Grill 2020. SPARC_Duke_Grill_OT2-OD025340_HumanVagusNerve_Claudin1IHC_Morphology. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.bh37j8rn>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Jun 30, 2020

Last Modified: Jun 30, 2020

PROTOCOL integer ID:
38751

SPARC_Duke_Grill_OT2-OD025340_HumanVagusNerve_Claudin1IHC_Morphology V.3

Nicole A. Pelot¹, J. Ashley Ezzell¹, Gabriel B. Goldhagen¹, Jake E. Cariello¹, Kara A. Clissold¹, Warren M. Grill¹

¹Duke University

SPARC

Tech. support email: info@neuinfo.org

Nicole A Pelot

Duke University

DISCLAIMER

DISCLAIMER – FOR INFORMATIONAL PURPOSES ONLY; USE AT YOUR OWN RISK

The protocol content here is for informational purposes only and does not constitute legal, medical, clinical, or safety advice, or otherwise; content added to protocols.io is not peer reviewed and may not have undergone a formal approval of any kind. Information presented in this protocol should not substitute for independent professional judgment, advice, diagnosis, or treatment. Any action you take or refrain from taking using or relying upon the information presented here is strictly at your own risk. You agree that neither the Company nor any of the authors, contributors, administrators, or anyone else associated with protocols.io, can be held responsible for your use of the information contained in or linked to this protocol or any of our Sites/Apps and Services.

ABSTRACT

The protocol describes immunohistochemistry with anti-claudin-1, imaging, image segmentation, and image analysis methods to quantify human vagus nerve morphology.

Keywords: Vagus nerve, nerve morphology, human vagus nerve, claudin-1, endoneurium, perineurium, epineurium, fascicles, image segmentation

MATERIALS

- Microscope slides with paraffin slices
- Xylene
- Ethanol
- Deionized water
- HIER Buffer L (Thermo, TA-135-HBL)
- H₂O₂
- Tris buffer
- Tris Tween buffer
- DAKO Protein Block (X0909)
- Antibody Diluent OP Quanto (Thermo, TA-125-ADQ)
- Rabbit anti-claudin-1 (Abcam, ab15098)
- Biotinylated SP-conjugated Affinipure goat anti-rabbit IgG (H+L) (Jackson, 111-065-144)
- ABC Elite (Vector, PK-6100)
- DAB chromogen (Thermo, TA-125-QHDX)
- Harris hematoxylin (Thermo, 6765003)
- DPX mountant (Electron Microscopy Sciences, 13512)
- Microscope with color camera
- Nikon's NIS Elements
- Matlab

Immunohistochemistry

- 1 Bake slides with sections of paraffin-embedded vagus nerve at 50°C overnight.
- 2 Deparaffinize the slides and hydrate them to distilled water: xylene (2x 6 min), 100% ethanol (5 min), 95% ethanol (4 min), 70% ethanol (3 min), deionized water (2x 1 min).
- 3 Perform heat-induced epitope retrieval (HIER) at 120°C for 30 s followed by 90°C for 10 s, using a buffer with pH 6.0 (Thermo, TA-135-HBL).
- 4 Cool for 20 min at room temperature.

- 5 Rinse in deionized water (2x 2 min).
- 6 Block with 3% H₂O₂ diluted in deionized water for 10 min.
- 7 Rinse in deionized water (2x 2 min).
- 8 Rinse in Tris buffer (1x 2 min).
- 9 Block using DAKO Protein Block (X0909) for 10 min at room temperature.
- 10 Apply the primary antibody (rabbit anti-claudin-1, Abcam, ab15098) diluted in Thermo Antibody Diluent to a concentration of 1:50, and incubate overnight at 4°C.
- 11 Rinse in Tris Tween buffer (2x 2 min).
- 12 Rinse in Tris buffer (1x 2 min).

- 13** Apply the secondary antibody (biotinylated SP-conjugated Affinipure goat anti-rabbit IgG (H+L), Jackson, 111-065-144) diluted in Thermo Antibody Diluent to a concentration of 1:500, and incubate for 1 hour at room temperature.
- 14** Rinse in Tris Tween buffer (2x 2 min).
- 15** Rinse in Tris buffer (1x 2 min).
- 16** Apply ABC Elite (Vector, PK-6100) at a concentration of 1:50 for 30 min at room temperature.
- 17** Rinse in Tris Tween buffer (2x 2 min).
- 18** Rinse in Tris buffer (1x 2 min).
- 19** Apply DAB chromogen (Thermo, TA-125-QHDX) for 1 min at room temperature.
- 20** Rinse in deionized water (2x 2 min).
- 21** Counterstain using hematoxylin.

- 22 Dehydrate, clear, and coverslip using DPX mountant.

Microscopy

- 23 Each sample was imaged at 10x using a Nikon Ti2 microscope with a Photometrics Prime 95B-25MM camera (Nikon Instruments Inc.). We selected the best of four slices for each sample based on the quality of the slice (no tearing or fraying).

Image Segmentation

- 24 We used Nikon's NIS Elements software (v5.02.01, Build 1270) to segment human vagus nerve immunohistochemical micrographs (anti-claudin-1) using the General Analysis RGB tool.
- 25 For each image, we selected preprocessing steps, such as smoothing and sharpening.
- 26 For each image, we selected ranges of hues, saturations, and intensities to values that identify the perineurium and different values to identify the entirety of the nerve.
- 27 For each image, we selected postprocessing steps, such as setting a minimum size criterion (eliminate small off-target regions), smoothing, cleaning, closing, and filling holes.
- 28 We made manual adjustments as needed, including manual deletion of off-target regions and filling of target areas that had not been captured.

- 29 We converted the binary segmented image into “Graticule Masks”, binary images saved as TIFs.

Image Analysis

- 30 We imported the TIFs into Matlab and generated a data structure of the x and y coordinates of the pixels for each closed boundary of the loaded binary images using the *bwboundaries* function.
- 31 We stored the pixel coordinates with indexing that assigns a fascicle number, which were then checked so that both the interior and exterior perineurium trace relate to the same fascicle.
- 32 We scaled the pixel coordinates to microns using the segmented scale bar.
- 33 We calculated cross-sectional area of each fascicle (inner perineurium and outer perineurium traces) and nerve using Matlab’s *polyarea*. Effective diameter (for a nerve or fascicle) is the diameter of the circle that has the same cross-sectional area as the raw trace. The perineurium thickness is half of the difference in effective diameters of the inner and outer perineurium traces.