JUL 09, 2020

ssDNA Extraktion

PLOS One

Franziska Patzold¹

¹Senckenberg Natural History Collections Dresden

Franziska Patzold

ABSTRACT

ssDNA Extraction Protocol

This protocol is based on the Oligonucleotide extraction protocol of the NEB PCR & Clean-up Kit and the Qiagen DNeasy Protocol with a few modifications to maximize DNA output. First results of this protocol will be published soon, including first sequencing results of the so extracted DNA. Please understand that this protocol is so far only tested for tissue (legs and abdomen) of invertebrates (Lepidoptera, genus *Hyles*). I recommend testing this protocol with tissue of non valuable specimens first.

GUIDELINES

Make sure to not touch the specimen/tissue at any given point, **always wear gloves** and **clean all used forcecepts before and after usage** with DNAaway or similar.

Make sure to clean the sample if possible before you start the Lysis (by UV-light or chlor, not possible in insect legs or soft tissue).

Only use this Extraction protocol in a cleaned environment, ideally in an aDNA-laboratory.

Take full pre-cautions;

- clean your whole bench before DNA extraction
- always clean your gloves by using DNAaway after touching the tube of a sample
- take care of aerosols! do not touch/open any tube while your gloves are still wet
- of cause only use filter-tips

MATERIALS

MATERIALS

Monarch® PCR & DNA Cleanup Kit (5 µg) **New England Biolabs Catalog #T1030**

Lysis Buffer FN **LGC biosearch Catalog #NAP41031**

Proteinase K **Analytik Jena Catalog #845-CH-0010006**

OPEN ACCESS

DOI:
[dx.doi.org/10.17504/protocols.io.8m9hu96](https://doi.org/10.17504/protocols.io.8m9hu96)

External link:
<https://doi.org/10.1371/journal.pone.0235222>

Protocol Citation: Franziska Patzold 2020. ssDNA Extraktion. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.8m9hu96>

MANUSCRIPT CITATION:
Patzold & Hundsdoerfer 2020, in preparation

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

Created: Oct 23, 2019

Last Modified: Jul 09, 2020

PROTOCOL integer ID:
29089

Keywords: ssDNA,
extraction, aDNA, museum

SAFETY WARNINGS

! don't drink the Ethanol ;)

BEFORE START INSTRUCTIONS

Input amount of DNA to be purified should not exceed the binding capacity of the column (5 µg). A starting sample volume of 50 µl is recommended. For smaller samples, nuclease-free water can be used to adjust the volume to the recommended volume range. Centrifugation should be carried out at 16,000 x *g* in a standard laboratory microcentrifuge at room temperature.

Buffer Preparation:

Add ethanol to Monarch DNA Wash Buffer prior to use (4 volumes of ≥ 95% ethanol per volume of Monarch DNA Wash Buffer).

- For 50-prep kit add 20 ml of ethanol to 5 ml of Monarch DNA Wash Buffer
- For 250-prep kit add 100 ml of ethanol to 25 ml of Monarch DNA Wash Buffer

Always keep all buffer bottles tightly closed when not in use.

All centrifugation steps should be carried out at 16,000 x *g*. (~13K RPM in a typical microcentrifuge).

This ensures all traces of buffer are eluted at each step.

Elution

- 1 Place dry tissue in 1.5 ml microcentrifuge tube and grind it, using an oscillating mill (MM 400, Retsch) for **1 min** at a frequency of **25 Hz**, using two sterile stainless steel balls with a diameter of 3 mm

Lysis

- 2 Insert to the ground tissue

🧴 45 µL Lysis buffer

🧴 5 µL Proteinase K

50 µL total volume mix well by handshaking, **don't vortex!** and gently spin down the solution.

- 3 Put the tube in an
Mixing Block (MB-102, Thermocell)

56 °C

08:00:00 at least, better overnight

Mix gently, and spin again.

Extraction

- 4 Add

100 µL DNA Cleanup Binding Buffer

to the

50 µL sample

mix by pipetting up and down.

- 5 Add

300 µL ethanol ($\geq 95\%$)

Mix well by pipetting up and down or flicking the tube. **Do not vortex.**

- 6 Insert column into collection tube,
load sample onto column and close the cap.

00:01:00 Centrifuge

then discard flow-through

- 7 Re-insert column into collection tube.

Add

500 µL DNA Wash Buffer

and

00:01:00 Centrifuge

Discard flow-through.

- 8 Repeat Step 7.

This step is recommended for removal of enzymes that may interfere with downstream applications (e.g., Proteinase K).

- 9** **Transfer column** to a clean 1.5 ml microfuge tube.
Use care to ensure that the tip of the column does not come into contact with the flow-through. If in doubt, re-spin for 1 minute to ensure traces of salt and ethanol are not carried over to the next step.

Two-Step-Elution

- 10** Add
17.5 µL Elution Buffer
to the center of the matrix
Wait for
00:10:00 Incubation
then spin for
00:01:00 Centrifuge
to elute the DNA.
- 11** Repeat Step 10 to extract all DNA