SEP 21, 2021

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.wxrfm6

Protocol Citation: Hongrong Yang, Junfei Xia, James Monaghan, Heather A Clark 2021. Staining and imaging of mouse submandibular ganglion by α -bungarotoxin and nanosensor. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.wxrfm6>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: Jan 09, 2019

Last Modified: Sep 21, 2021

PROTOCOL integer ID:
 19153

Staining and imaging of mouse submandibular ganglion by α -bungarotoxin and nanosensor

Hongrong Yang¹, Junfei Xia¹, James Monaghan¹, Heather A Clark¹

¹Northeastern University

SPARC

Tech. support email: info@neuinfo.org

Junfei Xia

ABSTRACT

This protocol describes how to label acetylcholine receptors in isolated submandibular ganglion from mice using α -bungarotoxin and nanosensor formulated by the Clark lab.

MATERIALS

MATERIALS

100% formalin **Merck MilliporeSigma (Sigma-Aldrich) Catalog #F1635-500ML**

α -Bungarotoxin Alexa Fluor 647 conjugated **Thermo Fisher Scientific Catalog #B35450**

PBS pH 7.4 **Thermo Fisher Scientific Catalog #10010023**

ChAT^{BAC}-eGFP transgenic mice from The Jackson Laboratory.
 Stock number: 007902

Extraction of submandibular ganglion

1. Enthanize animals using CO2.
2. Make a midline incision in the neck and expose the salivary glands.
3. Separate the pair of salivary glands and expose the salivary ducts.
4. Remove connective tissues surrounding salivary ducts and cut the ducts from where they enter salivary glands to where they enter digastric muscles. The submandibular ganglion are attached to cutted ducts.

Expected result

Well isolated salivary ducts with at least one submandibular ganglion attached

Staining of submandibular ganglion by α -bungarotoxin and ...

2. 1. Rinse submandibular ganglion with PBS for 00:20:00
2. Soak submandibular ganglion in 500 μ L 0.000002 Molarity (M) α -Bungarotoxin Alexa Fluor 647 conjugated Thermo Fisher Scientific Catalog #B35450 for 00:30:00 .
- Alternatively, soak submandibular ganglion in 500 μ L nanosensor solution formulated by Clark Lab for 00:30:00
3. Rinse submandibular ganglion with PBS for 00:20:00 .

Imaging of submandibular ganglion

3. 1. Place the submandibular ganglion on a glass slide and cover it with cover slip.
2. Image the whole tissue using a Zeiss LSM 700 confocal. Use 488 nm laser line for GFP and Oregon Green 488 channel, 555 nm laser line for pHAb, and 639 nm laser line for bungarotoxin-Alexa Fluor 647 channel. Set laser power at 2% or at the appropriate power intensity that cellular structures can be clearly seen.

Software

ZEN pro

NAME

ZEISS

DEVELOPER