OCT 14, 2019

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.73nhqme

Protocol Citation: Elinor Gottschalk, Bulent Arman Aksoy, Pinar Aksoy, Eric Czech, Jeff Hammerbacher 2019. Staining cells with IncuCyte Cytolight Rapid Dyes for flow cytometry or fluorescent microscopy.
protocols.io
<https://dx.doi.org/10.17504/protocols.io.73nhqme>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: Oct 09, 2019

Last Modified: Oct 14, 2019

PROTOCOL integer ID:
 28494

Staining cells with IncuCyte Cytolight Rapid Dyes for flow cytometry or fluorescent microscopy

Elinor Gottschalk¹, Bulent Arman Aksoy¹, Pinar Aksoy¹, Eric Czech¹, Jeff Hammerbacher¹

¹Medical University of South Carolina

Hammer Lab

Tech. support phone: +18437924527 email: arman@hammerlab.org

Elinor Gottschalk

ABSTRACT

This protocol is for labeling cells with IncuCyte Cytolight Rapid dyes from Essen Bioscience. We optimized the dye concentration for Jurkat cells, pmel-1 T cells, and B16F10 melanoma cells.

GUIDELINES

We optimized [IncuCyte® CytoLight Rapid Green Reagent for live cell cytoplasmic labeling](#) with Jurkat cells, pmel-1 T cells, and [IncuCyte® CytoLight Rapid Red Reagent](#) with B16F10 melanoma cells. The optimal concentrations (based on minimal decrease in cell viability and highest percentage of stained cells) were:

- Jurkat: 1 µM (green)
- pmel-1 T cells: 0.11 µM (green)
- B16F10 cells: 0.11 µM (red)

We visualized the stained cells by flow cytometry or on the Keyence BZ-X710 fluorescence microscope equipped with GFP and Cy5 filters.

MATERIALS

MATERIALS

IncuCyte® CytoLight Rapid Green Reagent for live cell cytoplasmic labeling **Essen Biosciences Catalog #4705**

1X PBS **VWR Scientific Catalog #75800-986**

IncuCyte® CytoLight Rapid Red Reagent for live cell cytoplasmic labeling **Essen Biosciences Catalog #4706**

1 Harvest cells and wash with 1X PBS

2 Centrifuge 350 x g , 5 minutes

2.1 Resuspend cells in PBS at 1 million cells per ml

3 Add IncuCyte Dye

- For pmel-1 T cells: we determined that 0.11 μ M of green dye was optimal
- We made our working stock 11 μ M (so for every 1 ml of cell suspension, we add 10 μ l of working stock of dye)
- For Jurkat cells, 1 μ M of green dye was optimal
- For B16F10 cells, 0.11 μ M of red dye was optimal

3.1 Incubate the cells at 37 °C for 00:20:00 (we wrapped the tube in foil and placed it in a water bath).

- invert the tube twice during incubation to mix the cells

3.2 Add a 6-fold volume of complete media (containing serum) to bind the excess dye.

3.3 Centrifuge 350 x g , 5 minutes

4 Resuspend cell in complete media at the desired concentration.

- Two different cell types can be labeled with different dyes and then co-cultured and visualized
- Stained T cells can be visualized migrating in collagen gel

5 Assay the cells via flowcytometry or fluorescence microscopy.
Flow cytometry: FITC for green dye; APC for red dye

Fluorescence microscopy: FITC filter for green dye; Cy5 filter for red dye