NOV 07, 2017

OPEN ACCESS

DOI:
[dx.doi.org/10.17504/protocols.io.ki6cuhe](https://doi.org/10.17504/protocols.io.ki6cuhe)

External link:
<https://doi.org/10.1371/journal.pone.0193657>

Protocol Citation: Sven Van Poucke, Dana Huskens, Kurt Van der Speeten, Mark Roest, Bart Lauwereins, Ming-Hua Zheng, Seppe Dehaene, Joris Penders, Abraham Marcus, Marcus Lancé 2017. Standard hematological analyses HIPEC . **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.ki6cuhe>

MANUSCRIPT CITATION:
 Poucke SV, Huskens D, Speeten KVd, Roest M, Lauwereins B, Zheng M, Dehaene S, Penders J, Marcus A, Lancé M (2018) Thrombin generation and platelet activation in cytoreductive surgery combined with hyperthermic intraperitoneal chemotherapy - A prospective cohort study. PLoS ONE 13(6): e0193657. doi: [10.1371/journal.pone.0193657](https://doi.org/10.1371/journal.pone.0193657)

Standard hematological analyses HIPEC

PLOS One

Sven Van Poucke¹, Dana Huskens², Kurt Van der Speeten³, Mark Roest², Bart Lauwereins⁴, Ming-Hua Zheng⁵, Seppe Dehaene⁴, Joris Penders⁶, Abraham Marcus⁷, Marcus Lancé⁷

¹Department of Anesthesiology, Intensive Care, Emergency Medicine and Pain Therapy, ZOL, Genk, Belgium;

²Synapse Research Institute, Maastricht, The Netherlands.;

³Department of Surgery, ZOL, Genk, Belgium.;

⁴Department of Anesthesiology, Intensive Care, Emergency Medicine and Pain Therapy, ZOL, Genk, Belgium.;

⁵Department of Hepatology, the First Affiliated Hospital of Wenzhou Medical University, Wenzhou 325000, China.;

⁶Central Diagnostic Laboratory, ZOL, Genk, Belgium.;

⁷Department of Anesthesiology, ICU and Perioperative Medicine, HMC, Doha, Qatar.

HIPACAT

Sven Van Poucke

Department of Anesthesiology, Intensive Care, Emergency Medi...

ABSTRACT

Standard hematological analyses
 EDTA-anticoagulated blood was used for cytometric analysis using a whole blood counter Sysmex XE 2100® (Sysmex, Kobe, Japan) to obtain a whole blood count. Fibrinogen levels were determined with an ACL-9000 (Diamond Diagnostics, Holliston, MA) coagulation analyser, using a PT-fibrinogen high sensitivity reagent, which is a high-sensitivity calcium thromboplastin that allows the simultaneous determination of PT and fibrinogen levels. aPTT was measured using an ACL-9000 coagulation analyser and INR was calculated by the formula INR (PT patient/PT normal).

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

Created: Nov 01, 2017

Last Modified: Mar 08, 2018

PROTOCOL integer ID:
8510

Keywords: HIPEC