

Jul 27, 2019

## Streaking bacteria on an agar plate from a glycerol stock

DOI

dx.doi.org/10.17504/protocols.io.5xig7ke

Cristian Riccio<sup>1</sup>

<sup>1</sup>University of Cambridge



**Cristian Riccio** 

University of Cambridge

#### Create & collaborate more with a free account

Edit and publish protocols, collaborate in communities, share insights through comments, and track progress with run records.

Create free account





DOI: https://dx.doi.org/10.17504/protocols.io.5xig7ke

**Protocol Citation:** Cristian Riccio 2019. Streaking bacteria on an agar plate from a glycerol stock. **protocols.io** <a href="https://dx.doi.org/10.17504/protocols.io.5xig7ke">https://dx.doi.org/10.17504/protocols.io.5xig7ke</a>

**License:** This is an open access protocol distributed under the terms of the **Creative Commons Attribution License**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: July 27, 2019



Last Modified: July 27, 2019

Protocol Integer ID: 26314

Keywords: bacteria on an agar plate, streaking bacteria, glycerol stock, agar plate, bacteria

### **Materials**

#### **MATERIALS**

Metal loop for microbiology

X Agar plate

**X** Bacterial glycerol stock

🛭 Bunsen burner

X Dry ice

# Troubleshooting



1 Transfer the glycerol stock onto dry ice next to a flame. 2 Sterilise the loop in the flame. 3 Rub the loop in the glycerol stock. 4 Streak the bacteri in zigzag over half the agar plate. 5 Sterilise the loop. 6 Zigzag over half the plate which overlaps with the first half by a quarter of a plate. 7 Sterilise the loop. 8 Zigzag over half the plate which overlaps with the second half by a quarter of a plate. 9 Incube overnight at the temperature which is optimal for bacterial growth, 37 C for E. coli. 10 The next day, you should have single colonies visible on a quarter of the plate or half the plate.