JUL 27, 2019

Streaking bacteria on an agar plate from a glycerol stock

Cristian
Riccio¹

¹University of Cambridge

Cristian Riccio
University of Cambridge

MATERIALS

MATERIALS

⊗ Metal loop for microbiology Contributed by users

⊗ Agar plate Contributed by users

⊗ Bacterial glycerol stock Contributed by users

⊗ Bunsen burner Contributed by users

⊗ Dry ice Contributed by users

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.5xig7ke

Protocol Citation: Cristian Riccio 2019. Streaking bacteria on an agar plate from a glycerol stock.
protocols.io
<https://dx.doi.org/10.17504/protocols.io.5xig7ke>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Jul 27, 2019

Last Modified: Jul 27, 2019

PROTOCOL integer ID:
26314

- 1 Transfer the glycerol stock onto dry ice next to a flame.

- 2 Sterilise the loop in the flame.
- 3 Rub the loop in the glycerol stock.
- 4 Streak the bacteri in zigzag over half the agar plate.
- 5 Sterilise the loop.
- 6 Zigzag over half the plate which overlaps with the first half by a quarter of a plate.
- 7 Sterilise the loop.
- 8 Zigzag over half the plate which overlaps with the second half by a quarter of a plate.
- 9 Incube overnight at the temperature which is optimal for bacterial growth, 37 C for *E. coli*.

10 The next day, you should have single colonies visible on a quarter of the plate or half the plate.