JUL 31, 2018

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.qxfdxjn

External link:
<https://doi.org/10.1371/journal.pone.0201432>

Protocol Citation: Marie Zikanova 2018. Study of purinosome assembly in cell-based model systems with de novo purine synthesis and salvage pathway deficiencies. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.qxfdxjn>

MANUSCRIPT CITATION:
 Baresova V, Skopova V, Souckova O, Krijt M, Kmoch S, Zikanova M (2018) Study of purinosome assembly in cell-based model systems with *de novo* purine synthesis and salvage pathway deficiencies. PLoS ONE 13(7): e0201432. doi: [10.1371/journal.pone.0201432](https://doi.org/10.1371/journal.pone.0201432)

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Study of purinosome assembly in cell-based model systems with de novo purine synthesis and salvage pathway deficiencies

PLOS One

Marie Zikanova¹

¹Research Unit for Rare Diseases, Department of Paediatrics and Adolescent Medicine, First Faculty of Medicine, Charles University and General University Hospital, Prague, Czech Republic

Marie Zikanova

ABSTRACT

The purinosome has been observed in a broad spectrum of cells, but some studies claim that it is an artefact of the constructs used for visualization or stress granules resulting from the exposure of cells to nutrient-reduced growth media. Both may be true depending on the method of observation. To clarify this point, we combined two previously used methods, transfection and immunofluorescence, to detect purinosomes in purinosome-free cells deficient in particular DNPS steps (CR-DNPS cells) and in cells deficient in the salvage pathway, which resulted in construction of the purinosome regardless of purine level (CR-HGPRT cells). To restore or disrupt purinosome formation, we transiently transfected CR-DNPS and CR-HGPRT cells with vectors encoding BFP-labelled wild-type (wt) proteins and observed the normalization of purinosome formation. The CR-DNPS cell line transfected with a DNA plasmid encoding an enzyme with zero activity served as a negative control for purinosome formation.

ATTACHMENTS

[materials_for_io.docx](#)

Protocol status: Working

Created: Jun 13, 2018

Last Modified: Jul 31, 2018

PROTOCOL integer ID:
12999

Keywords: purinosome,
transfection,
immunofluorescence, CRISPR

1