

Mar 22, 2024

# Study Protocol for the AIRCARD Study: A Prospective Cohort Study Utilizing DANCAVAS and VIVA Screening Trials

DOI

[dx.doi.org/10.17504/protocols.io.kxygxykqzl8j/v1](https://dx.doi.org/10.17504/protocols.io.kxygxykqzl8j/v1)

Stephan Peronard Mayntz<sup>1,2,3</sup>, Roda Abdulkadir Mohamed<sup>1,2,3</sup>, Anna Mejldal<sup>1</sup>, Jens-Jakob Kjer Møller<sup>1</sup>, Jes Lindholt<sup>1,4</sup>, Axel Diederichsen<sup>1,5</sup>, Lise Marie Frohn<sup>6</sup>, Jørgen Brandt<sup>6</sup>, Matthias Ketzel<sup>6</sup>, Jibrán Khan<sup>6</sup>, Jess Lambrechtsen<sup>1,2,3</sup>

<sup>1</sup>OPEN – Open Patient data Explorative Network, Odense University Hospital, Region of Southern Denmark;

<sup>2</sup>Department of Clinical Research, University of Southern Denmark;

<sup>3</sup>Cardiology Research Unit, Odense University Hospital, Svendborg;

<sup>4</sup>Department of Cardiothoracic and Vascular Surgery, Odense University Hospital, Odense, Denmark;

<sup>5</sup>Department of Cardiology, Odense University Hospital, Odense, Denmark;

<sup>6</sup>Department of Environmental Science, Aarhus University, Roskilde

Stephan Peronard Mayntz: +45 42 66 32 23;



**Stephan Peronard Mayntz**

Odense University Hospital

## Create & collaborate more with a free account

Edit and publish protocols, collaborate in communities, share insights through comments, and track progress with run records.

Create free account

OPEN  ACCESS



**DOI:** <https://dx.doi.org/10.17504/protocols.io.kxygxykqzl8j/v1>

**Protocol Citation:** Stephan Peronard Mayntz, Roda Abdulkadir Mohamed, Anna Mejldal, Jens-Jakob Kjer Møller, Jes Lindholt, Axel Diederichsen, Lise Marie Frohn, Jørgen Brandt, Matthias Ketzel, Jibran Khan, Jess Lambrechtsen 2024. Study Protocol for the AIRCARD Study: A Prospective Cohort Study Utilizing DANCAVAS and VIVA Screening Trials. **protocols.io**  
<https://dx.doi.org/10.17504/protocols.io.kxygxykqzl8j/v1>

**License:** This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

**Protocol status:** Working

**We use this protocol and it's working**

**Created:** March 18, 2024

**Last Modified:** March 22, 2024

**Protocol Integer ID:** 97132

**Keywords:** noise pollution exposure on cardiovascular disease, noise pollution on cardiovascular disease, study protocol for the aircard study, cardiovascular disease burden, noise pollution exposure, term exposure to air pollutant, cardiovascular disease incidence, aircard study, prospective cohort study utilizing dancava, term exposure to air, noise pollution through the nord2000, including traditional cardiovascular risk factor, cardiovascular health, noise pollution, differential impacts of air, pollution exposure, robust prospective cohort analysis, prospective cohort study, traditional cardiovascular risk factor, air pollutant, viva screening trials background, cox proportional hazards model, cardiovascular disease, public health measure, need for public health measure, heart disease, proportional hazards model, family history of heart disease, pollution, pollutant, mortality, cohort, smoking status, airgis modelling system

## Abstract

**Background:** The AIRCARD study is designed to investigate the relationship between long-term exposure to air and noise pollution and cardiovascular disease incidence and mortality.

**Aim:** To conduct a robust prospective cohort analysis assessing the cumulative and differential impacts of air and noise pollution exposure on cardiovascular disease and mortality. This study will adjust for relevant confounders, including traditional cardiovascular risk factors, socioeconomic indicators, and medication use.

**Methods:** This prospective cohort study will include male participants aged 65-74, recruited from the two large Danish DANCAVAS and VIVA trials, both population-based randomized, multicentered, clinically controlled studies. We will assess long-term exposure to air pollutants using the state-of-the-art DEHM/UBM/AirGIS modelling system and noise pollution through the Nord2000 and SoundPLAN models, covering data from 1979 to 2019. The primary analysis will utilize Cox proportional hazards models, adjusted for confounders identified in the cohort (age, body mass index, hypertension, diabetes, smoking status, family history of heart disease, socioeconomic factors, and medication use).

**Discussion:** The AIRCARD study will address global concerns about the impact of air and noise pollution on cardiovascular disease. This research is crucial for understanding how the pollutants contribute to cardiovascular disease. We aim to provide valuable insights into this area, emphasizing the need for public health measures to mitigate pollution exposure. Our goal is to provide policymakers and healthcare professionals with information on the role of environmental factors in cardiovascular health that could influence global strategies to reduce the cardiovascular disease burden associated with pollution.

## Attachments



Study Protocol for t...

98KB



## Guidelines

### **Ethics approval and consent:**

The DANCAS and VIVA studies were approved from "Videnskabsetisk Komité" (S20140028, S20160164 and M20080028). Access to the participants' former residential addresses has been approved from "Sundhedsdatastyrelsen" (FSEID-00005213).

### **Feasibility:**

We have formed the group of researchers that will perform the study and we have received all the necessary approvals. The group includes professors in the areas where special skills are needed.

### **COLLABORATION:**

This project is a collaboration across disciplines and sectors (cardiology, physics, environmental sciences, surgery, public health). The study will be performed as a collaboration between:

- Cardiovascular Research Unit, Odense University Hospital - Svendborg; Professor Jess Lambrechtsen, Professor Kenneth Egstrup, Stephan Peronard Mayntz, MD, MPH, and Roda A. Mohamed, MD.
- Department of Environmental Science, Aarhus University, Roskilde; Associate Professor Lise M. Frohn, Professor Jørgen Brandt, Associate Professor Matthias Ketznel, and Postdoctoral Fellow Jibran Khan.
- Department of Cardiology, Odense University Hospital; Professor Axel Cosmus Pyndt Diederichsen.
- Department of Cardiac, Thoracic and Vascular Surgery, Odense University Hospital; Professor Jes Sanddal Lindholt.
- OPEN – Open Patient data Explorative Network, Odense University Hospital, Region of Southern Denmark; Anna Mejlidal, Jens-Jakob Kjer Møller.

### **Funding:**

Supported by the Region of Southern Denmark and by Shipowner Per Henriksen's Foundation.

### **Availability of data and materials:**

A Data Management Plan and Sharing Statement has been developed and is publicly available at [dmp.deic.dk](https://dmp.deic.dk) with ID: 6570

## Troubleshooting

## Project design

- 1 This study will be designed as a prospective registry-based observational cohort study using modelled air and noise pollution data. The population is males from two Danish clinical trials (DANCAVAS and VIVA trials).

## Study population

- 2 DANCAVAS (39-41) was a population-based randomized, multicentered, clinically controlled studies designed to evaluate the benefits of 7-step multiple cardiovascular screening and modern vascular prophylaxis in a population of men, aged 65-74 years, living in the southern part of Denmark.
- 3 For each participant, all relevant CVD risk factors were measured or determined.  
  
The screening included:
  1. Low-dose non-contrast CT scan to detect CAC and aortic/iliac aneurysms
  2. Ankle-brachial blood pressure index (ABI) to detect peripheral arterial disease (PAD) and hypertension
  3. A telemetric assessment of the heart rhythm, and
  4. A measurement of the cholesterol and plasma glucose levels.
- 4 The Viborg Vascular (VIVA) screening trial (42) was a randomized, multicentered, population-based clinically controlled study designed to evaluate the benefits of vascular screening and modern vascular prophylaxis in men between 65-74 years of age living in the region of Mid Denmark (Viborg County).
  - For each participant, abdominal ultrasound scanning of the infrarenal aorta were performed to detect abdominal aortic aneurism (AAA), and ABI were measured to detect PAD and hypertension (43).
- 5 In both trials, an AAA was defined as maximal infrarenal diameter of 30 mm or more, and PAD was defined as an ABI < 0.90 or ≥1.40 using the same validated hand held Doppler-based methodology (44).
  - We have accounted for traditional CVD confounders in all participants. This is unique for our study.
  - Previous studies on air pollution and CVD burden have not been able to prospectively consider these confounding variables to the same extent.

## Danish national registries

- 6 The study will use Danish National Registries through Statistics Denmark and the Danish Health Data Authority.
- 7 These registries include the Cause of Death Register, the National Patient Register, the Civil Registration System, the Income Statistics Register, the Danish Education Registers, the Family Income Register, the National Prescription Registry, and the Employment Classification Module.
- 8 Each registry offers unique and comprehensive data, from mortality statistics to socioeconomic variables, contributing to a multifaceted understanding of the impacts of pollution on CVD. The participants were followed in the registries from 1979 to 2019.

## Air pollution

- 9 In Denmark a validated and reliable air pollution model system is available. The system is named DEHM/UBM/AirGIS (28, 30–32) and consists of three coupled models; the Danish Eulerian Hemispheric Model (DEHM) (28, 32), the Urban Background Model (UBM) (29) and the Operational Street Pollution Model (OSPM) (30, 31) and a GIS system (AirGIS) that couples the modelled concentrations with the address level of the population.
- 10 The system calculates air pollution concentrations of 80 chemical species as well as air pollution levels in cities, in streets and on address level even on both side of the street. These pollution levels can be calculated back to 1979 giving data 40 years back.
- 11 The model system is validated in relation to air pollution measurements throughout Denmark back to 1990 with high correlation between model estimated values and measured values (28, 30, 31, 45). This multi-scale model system is unique, capable of running on very high temporal (hourly) and spatial (address level) resolutions.
- 12 The development of the models and the calculation of air pollution and measurements is performed at Aarhus University, Department of Environmental Science (ENVS). The model is robust; taking all necessary factors into account that could contribute to the individual life-long air pollution exposure and the model is one of the best in the world.

## Noise pollution

- 13 Noise is modelled using state-of-the-art algorithms implemented in a well-known software, the SoundPLAN.
- 14 The algorithms, reflecting advanced physics and mathematics-based knowledge, consider the propagation of sound in the atmosphere as well as the sound originating from the source, e.g., road transport, railway. Here, information from various national registers, such as the national traffic database, including traffic counts, travel speeds, the building register, the address register, and the Danish surface and elevation model.



- 15 In addition, advanced weather classes reflecting all meteorological conditions in Denmark are used in noise calculations. The model output is a noise estimate at the address location or any location of interest in Denmark, which can be subsequently used to investigate the health impacts of short-term and/or long-term noise exposure (46, 47).

## Statistics

- 16 The entire study population is monitored until December 31. 2019, in the Danish national registers.  
Primarily a multivariate Cox proportional hazards regression model will be used to examine the associations between air- and noise pollution and CVD morbidity and mortality when adjusting for inclusion date, sex, and other potential confounding factors at baseline.
- 17 A Statistical Analysis Plan (SAP) is underway and is currently being revised. We refer to the SAP for all relevant statistics.

## Protocol references

1. Manisalidis I, Stavropoulou E, Stavropoulos A, Bezirtzoglou E. Environmental and Health Impacts of Air Pollution: A Review. *Front Public Health*. 2020;8:14.
2. Shah AS, Langrish JP, Nair H, McAllister DA, Hunter AL, Donaldson K, et al. Global association of air pollution and heart failure: a systematic review and meta-analysis. *Lancet*. 2013;382(9897):1039-48.
3. Hayes RB, Lim C, Zhang Y, Cromar K, Shao Y, Reynolds HR, et al. PM2.5 air pollution and cause-specific cardiovascular disease mortality. *Int J Epidemiol*. 2020;49(1):25-35.
4. Mannucci PM. Air pollution levels and cardiovascular health: Low is not enough. *Eur J Prev Cardiol*. 2017;24(17):1851-3.
5. Shah AS, Lee KK, McAllister DA, Hunter A, Nair H, Whiteley W, et al. Short term exposure to air pollution and stroke: systematic review and meta-analysis. *Bmj*. 2015;350:h1295.
6. Thomson H, Thomas S, Sellstrom E, Petticrew M. Housing improvements for health and associated socio-economic outcomes. *Cochrane Database Syst Rev*. 2013(2):Cd008657.
7. Roger VL, Go AS, Lloyd-Jones DM, Benjamin EJ, Berry JD, Borden WB, et al. Executive summary: heart disease and stroke statistics--2012 update: a report from the American Heart Association. *Circulation*. 2012;125(1):188-97.
8. Lloyd-Jones DM, Leip EP, Larson MG, D'Agostino RB, Beiser A, Wilson PW, et al. Prediction of lifetime risk for cardiovascular disease by risk factor burden at 50 years of age. *Circulation*. 2006;113(6):791-8.
9. Murray CJ, Lopez AD. Alternative projections of mortality and disability by cause 1990-2020: Global Burden of Disease Study. *Lancet*. 1997;349(9064):1498-504.
10. Blaha MJ, Cainzos-Achirica M, Dardari Z, Blankstein R, Shaw LJ, Rozanski A, et al. All-cause and cause-specific mortality in individuals with zero and minimal coronary artery calcium: A long-term, competing risk analysis in the Coronary Artery Calcium Consortium. *Atherosclerosis*. 2020;294:72-9.
11. Budoff MJ, Mohlenkamp S, McClelland R, Delaney JA, Bauer M, Jockel HK, et al. A comparison of outcomes with coronary artery calcium scanning in unselected populations: the Multi-Ethnic Study of Atherosclerosis (MESA) and Heinz Nixdorf RECALL study (HNR). *J Cardiovasc Comput Tomogr*. 2013;7(3):182-91.
12. Diederichsen AC, Sand NP, Norgaard B, Lambrechtsen J, Jensen JM, Munkholm H, et al. Discrepancy between coronary artery calcium score and HeartScore in middle-aged Danes: the DanRisk study. *Eur J Prev Cardiol*. 2012;19(3):558-64.
13. Folsom AR, Kronmal RA, Detrano RC, O'Leary DH, Bild DE, Bluemke DA, et al. Coronary artery calcification compared with carotid intima-media thickness in the prediction of cardiovascular disease incidence: the Multi-Ethnic Study of Atherosclerosis (MESA). *Arch Intern Med*. 2008;168(12):1333-9.
14. Diderichsen F, Andersen I, Manuel C. Uilighed i sundhed - årsager og indsatser. København: Sundhedsstyrelsen. 2011.
15. Flachs EM, Eriksen L, Koch MB, Ryd JT, Dibba E, Skov-Ettrup L, et al. Sygdomsbyrden i Danmark - sygdomme. København: Sundhedsstyrelsen. 2015.
16. Niiranen TJ, Vasan RS. Epidemiology of cardiovascular disease: recent novel outlooks on risk factors and clinical approaches. *Expert Rev Cardiovasc Ther*. 2016;14(7):855-69.
17. Lambrechtsen J, Gerke O, Egstrup K, Sand NP, Norgaard BL, Petersen H, et al. The relation between coronary artery calcification in asymptomatic subjects and both traditional risk factors and living in the city centre: a DanRisk substudy. *J Intern Med*. 2012;271(5):444-50.
18. Geels C, Andersson C, Hanninen O, Lanso AS, Schwarze PE, Skjoth CA, et al. Future premature mortality due to O3, secondary inorganic aerosols and primary PM in Europe--sensitivity to changes in climate, anthropogenic



- emissions, population and building stock. *Int J Environ Res Public Health*. 2015;12(3):2837-69.
19. Hanninen O, Knol AB, Jantunen M, Lim TA, Conrad A, Rappolder M, et al. Environmental burden of disease in Europe: assessing nine risk factors in six countries. *Environ Health Perspect*. 2014;122(5):439-46.
20. Lelieveld J, Klingmuller K, Pozzer A, Poschl U, Fnais M, Daiber A, et al. Cardiovascular disease burden from ambient air pollution in Europe reassessed using novel hazard ratio functions. *Eur Heart J*. 2019;40(20):1590-6.
21. Auchincloss AH, Diez Roux AV, Dvorchak JT, Brown PL, Barr RG, Davignus ML, et al. Associations between recent exposure to ambient fine particulate matter and blood pressure in the Multi-ethnic Study of Atherosclerosis (MESA). *Environ Health Perspect*. 2008;116(4):486-91.
22. Bauer M, Moebus S, Mohlenkamp S, Dragano N, Nonnemacher M, Fuchsluger M, et al. Urban particulate matter air pollution is associated with subclinical atherosclerosis: results from the HNR (Heinz Nixdorf Recall) study. *J Am Coll Cardiol*. 2010;56(22):1803-8.
23. Gill EA, Curl CL, Adar SD, Allen RW, Auchincloss AH, O'Neill MS, et al. Air pollution and cardiovascular disease in the Multi-Ethnic Study of Atherosclerosis. *Prog Cardiovasc Dis*. 2011;53(5):353-60.
24. O'Neill MS, Diez-Roux AV, Auchincloss AH, Shen M, Lima JA, Polak JF, et al. Long-term exposure to airborne particles and arterial stiffness: the Multi-Ethnic Study of Atherosclerosis (MESA). *Environ Health Perspect*. 2011;119(6):844-51.
25. Kaufman JD, Adar SD, Allen RW, Barr RG, Budoff MJ, Burke GL, et al. Prospective study of particulate air pollution exposures, subclinical atherosclerosis, and clinical cardiovascular disease: The Multi-Ethnic Study of Atherosclerosis and Air Pollution (MESA Air). *Am J Epidemiol*. 2012;176(9):825-37.
26. Kaufman JD, Adar SD, Barr RG, Budoff MJ, Burke GL, Curl CL, et al. Association between air pollution and coronary artery calcification within six metropolitan areas in the USA (the Multi-Ethnic Study of Atherosclerosis and Air Pollution): a longitudinal cohort study. *Lancet*. 2017;388:696-704.
27. Hoffmann B, Moebus S, Mohlenkamp S, Stang A, Lehmann N, Dragano N, et al. Residential exposure to traffic is associated with coronary atherosclerosis. *Circulation*. 2007;116(5):489-96.
28. Frohn LM, Christensen J. H., Brandt J. Development of a High-Resolution Nested Air Pollution Model: The Numerical Approach. *Journal of Computational Physics*. 2002;179(1):68-94.
29. Brandt J, Christensen JH, Frohn LM, Berkowicz R. Air pollution forecasting from regional to urban street scale—implementation and validation for two cities in Denmark. *Physics and Chemistry of the Earth, Parts A/B/C*. 2003;28(8):335-44.
30. Hvidtfeldt U, Ketzel M, Sørensen M, Hertel O, Khan J, Brandt J, et al. Evaluation of the Danish AirGIS air pollution modeling system against measured concentrations of PM<sub>2.5</sub>, PM<sub>10</sub>, and black carbon. *Environmental Epidemiology*. 2018;2(2):1-11.
31. Khan J, Kakosimos K, Raaschou-Nielsen O, Brandt J, Jensen SM, Ellermann T, et al. Development and performance evaluation of new AirGIS – A GIS based air pollution and human exposure modelling system. *Atmospheric Environment*. 2019;198:102-21.
32. Christensen JH. The Danish eulerian hemispheric model — a three-dimensional air pollution model used for the arctic. *Atmospheric Environment*. 1997;31(24):4169-91.
33. Holst G, Thygesen M, Pedersen CB, Peel RG, Brandt J, Christensen JH, et al. Ammonia, ammonium, and the risk of asthma. *Environmental Epidemiology*. 2018;2(3).
34. Horsdal HT, Agerbo E, McGrath JJ, Vilhjalmsen BJ, Antonsen S, Closter AM, et al. Association of Childhood Exposure to Nitrogen Dioxide and Polygenic Risk Score for Schizophrenia With the Risk of Developing Schizophrenia. *JAMA Netw Open*. 2019;2(11):e1914401.

35. Brook RD, Newby DE, Rajagopalan S. Air Pollution and Cardiometabolic Disease: An Update and Call for Clinical Trials. *Am J Hypertens*. 2017;31(1):1-10.
36. Brauer M, Casadei B, Harrington RA, Kovacs R, Sliwa K. Taking a Stand Against Air Pollution-The Impact on Cardiovascular Disease: A Joint Opinion From the World Heart Federation, American College of Cardiology, American Heart Association, and the European Society of Cardiology. *Circulation*. 2021;143(14):e800-e4.
37. Global burden of 87 risk factors in 204 countries and territories, 1990-2019: a systematic analysis for the Global Burden of Disease Study 2019. *Lancet*. 2020;396(10258):1223-49.
38. Lambrechtsen J, Mayntz SK, Engdam KB, Egstrup K, Nielsen J, Steffensen FH, et al. Relation between Accumulated Air Pollution Exposure and Sub-Clinical Cardiovascular Disease in 33,723 Danish 60-74-Year-Old Males from the Background Population (AIR-CARD): A Method Article. *Cardiology*. 2021;146(1):19-26.
39. Diederichsen AC, Rasmussen LM, Sogaard R, Lambrechtsen J, Steffensen FH, Frost L, et al. The Danish Cardiovascular Screening Trial (DANCAVAS): study protocol for a randomized controlled trial. *Trials*. 2015;16:554.
40. Danish Cardiovascular Screening Trial II (DANCAVAS-II). <https://clinicaltrials.gov/ct2/show/NCT03946410>.
41. Kvist TV, Lindholt JS, Rasmussen LM, Sogaard R, Lambrechtsen J, Steffensen FH, et al. The DanCavas Pilot Study of Multifaceted Screening for Subclinical Cardiovascular Disease in Men and Women Aged 65-74 Years. *Eur J Vasc Endovasc Surg*. 2017;53(1):123-31.
42. Lindholt JS, Sogaard R. Population screening and intervention for vascular disease in Danish men (VIVA): a randomised controlled trial. *Lancet*. 2017;390(10109):2256-65.
43. Grondal N, Bramsen MB, Thomsen MD, Rasmussen CB, Lindholt JS. The cardiac cycle is a major contributor to variability in size measurements of abdominal aortic aneurysms by ultrasound. *Eur J Vasc Endovasc Surg*. 2012;43(1):30-3.
44. Joensen JB, Juul S, Abrahamsen J, Henneberg EW, Lindholt JS. Doppler ultrasound compared with strain gauge for measurement of systolic ankle blood pressure. *Angiology*. 2008;59(3):296-300.
45. Ketzal M, Brandt J, Ellermann T, Olesen HR, Berkowicz R, Hertel O. Evaluation of the Street Pollution Model OSPM for Measurements at 12 Streets Stations Using a Newly Developed and Freely Available Evaluation Tool. *Journal of Civil and Environmental Engineering*. 2012.
46. Khan J, Ketzal M, Jensen SS, Gulliver J, Thysell E, Hertel O. Comparison of Road Traffic Noise prediction models: CNOSSOS-EU, Nord2000 and TRANEX. *Environ Pollut*. 2021;270:116240.
47. Khan J, Kakosimos K, Jensen SS, Hertel O, Sørensen M, Gulliver J, et al. The spatial relationship between traffic-related air pollution and noise in two Danish cities: Implications for health-related studies. *Sci Total Environ*. 2020;726:138577.