



Mar 06, 2020

Superoxide Radical-scavenging Activity

 [PLOS One](#)

DOI

<https://dx.doi.org/10.17504/protocols.io.bdaxi2fn>

Jing Xu¹

¹Hainan University



Jing Xu

Hainan University

Create & collaborate more with a free account

Edit and publish protocols, collaborate in communities, share insights through comments, and track progress with run records.

Create free account

OPEN  ACCESS



DOI: <https://dx.doi.org/10.17504/protocols.io.bdaxi2fn>

External link: <https://doi.org/10.1371/journal.pone.0234435>

Protocol Citation: Jing Xu 2020. Superoxide Radical-scavenging Activity. **protocols.io**

<https://dx.doi.org/10.17504/protocols.io.bdaxi2fn>

Manuscript citation:

Zhou J, Yang Q, Zhu X, Lin T, Hao D, Xu J (2020) Antioxidant activities of *Clerodendrum cyrtophyllum* Turcz leaf extracts and their major components. PLoS ONE 15(6): e0234435. doi: [10.1371/journal.pone.0234435](https://doi.org/10.1371/journal.pone.0234435)



License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: March 06, 2020

Last Modified: March 06, 2020

Protocol Integer ID: 33847

Keywords: superoxide radical scavenging effect, scavenging activity the superoxide radical scavenging effect, superoxide radical, scavenging activity, scavenging, extract

Abstract

The superoxide radical scavenging effects were examined [11]. Briefly, extract (1 ml) was added to 1 ml of 50 μ M NBT solution and 1 ml of 468 μ MNADH, and a 1ml aliquot of 60 μ M PMS reaction mixture. After 5min, absorbance was read at 560 nm. The inhibition percent and IC₅₀ valueswere calculated.

