OCT 18, 2018

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.uteewje

Protocol Citation: Priota Islam 2018. Synchronization via Bleaching (Spot Bleaching) protocol. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.uteewje>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: Oct 18, 2018

Last Modified: Oct 18, 2018

PROTOCOL integer ID:
 16966

Synchronization via Bleaching (Spot Bleaching) protocol

Priota Islam¹

¹Imperial College London

Behavioural Genomics

Priota Islam
 Imperial College London

ABSTRACT

If you only need a small number of semi-synchronized worms or if you simply need to remove yeast or bacterial contaminants, this protocol is easier and less time-consuming than the full-scale egg prep. This procedure is also called 'spot bleaching'.

MATERIALS

MATERIALS

Sodium Hydroxide **Fisher Scientific Catalog #BP359500**

Sodium Hypochlorite Solution **Contributed by users**

2X Bleach solution:

Sodium hypochlorite solution - 4ml

Millipore water - 3.5 ml

NaOH solution - 2.5 ml

TOTAL - 10 ml

- 1 Pipette ~20µL of the 2X bleach solution onto the unseeded portion of a new plate.

- 2 Pick approximately 10 gravid adults from the contaminated plate with a metal pick and place into the bleach solution on the clean plate. Try to avoid bringing too much bacteria with the worms. (It is best to pick twice to ensure a good number of gravid is picked)
- 3 After about 10 minutes, the worms should break open and release the bleach resistant eggs.
- 4 Leave the plate right side up overnight at 20° C.
- 5 The next day, check to see that L1s have hatched
- 6 Pick about 10 L1s onto a new seeded plate and continue growing them as desired.