T Cell Activation with anti-CD3 Antibodies Protocol - Human V.3

Kelsey Knight¹

¹BioLegend

Sam Li
BioLegend

DOI:
[dx.doi.org/10.17504/protocol.s.io.tkbeksn](https://doi.org/10.17504/protocol.s.io.tkbeksn)

External link:
<https://www.biolegend.com/protocols/t-cell-activation-with-anti-cd3-antibodies-protocol-human/4271/>

Protocol Citation: Kelsey Knight . T Cell Activation with anti-CD3 Antibodies Protocol - Human. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.tkbeksn>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Created: Sep 13, 2018

Last Modified: Aug 29, 2019

PROTOCOL integer ID:
15715

MATERIALS

STEP MATERIALS

96-well flat-bottom tissue culture plates with lids **Corning Catalog #3596**

- Sterile PBS
- Anti-human CD3 Antibody
 - Clone UCHT1 (LEAF™ format, Cat. No. 300413/300414/300432; Ultra-LEAF™ format, Cat. No. 300437/300438)
 - Clone OKT3 (LEAF™ format, Cat. No. 317303/317304/317315; Ultra-LEAF™ format, Cat. No. 317325/317326)
 - Clone HIT3a (LEAF™ format, Cat. No. 300313/300314; Ultra-LEAF™ format, Cat. No. 300331/300332)
- Cell culture medium (e.g., RPMI-1640 or IMDM supplemented with 10% FBS and 2mM L-glutamine)
- Sterile single-cell suspension of Ficoll-Hypaque-purified peripheral blood mononuclear cells, isolated T cells, or T cell subsets
- 96-well flat-bottom tissue culture plates with lids (e.g., Costar® Cat. No. 3596)
- * Soluble forms of LEAF™ purified UCHT1 (1 µg/ml) or LEAF™ purified HIT3a (0.01 – 0.1 µg/ml) may be used to activate T cells from PBMC cell populations.

1 Prepare a 10 µg/ml solution of anti-CD3 (clone UCHT1, OKT3, or HIT3a) in sterile PBS.

2 Dispense 50 µl of the antibody solution to each microwell of the 96-well assay plate. For the unstimulated control wells, add 50 µl of sterile PBS.

96-well flat-bottom tissue culture plates with lids **Corning Catalog #3596**

3 Seal plate. Incubate at 37°C for 2 hours or 4°C overnight.

4 Aseptically decant antibody solution from the microwell plate.

5 Wash plate microwells 3 times with sterile PBS. Discard liquid.

6 Prepare single cell suspension of cells of interest in supplemented cell culture medium to $1-2 \times 10^6/\text{ml}$.

7 Aliquot 200 μl cell suspension into plate microwells. Cover with lid. Incubate at 37°C in 5% CO_2 and 100% humidity for 3 days.