Th1 Polarization of Mouse CD4+ Cells V.2

Kelsey Knight¹

¹BioLegend

Sam Li
BioLegend

GUIDELINES

DOI:

[dx.doi.org/10.17504/protocols.io.tncemaw](https://doi.org/10.17504/protocols.io.tncemaw)

External link:

<https://www.biolegend.com/protocols/th1-polarization-of-mouse-cd4--cells-protocol/4242/>

Protocol Citation: Kelsey Knight . Th1 Polarization of Mouse CD4+ Cells.

protocols.io

<https://dx.doi.org/10.17504/protocols.io.tncemaw>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Created: Sep 14, 2018

Last Modified: Aug 29, 2019

PROTOCOL integer ID:
15780

Reagent List:

- Sterile PBS
- Cell culture medium (RPMI 1640 supplemented with 10% FBS)
- Sterile 12-well plate
- Sterile 6-well plate
- RBC Lysis Buffer (Cat. No. 420301)
- Anti-mouse CD3ε, clone 145-2C11 (LEAF™ format, Cat. No. 100314)
- Anti-mouse CD28, clone 37.51, (LEAF™ format, Cat. No. 102112)
- Anti-mouse IL-4, clone 11B11, (LEAF™ format, Cat. No. 504108)
- Recombinant mouse IL-2 (carrier-free) (Cat. No. 575402)
- Recombinant mouse IL-12 (p70) (carrier-free) (Cat. No. 577002)
- Monensin Solution (Cat. No. 420701)
- PMA (Phorbol 12-myristate 13-acetate) (Cat. No. P8139 from Sigma)
- Ionomycin (Cat. No. I0634 from Sigma)

MATERIALS

MATERIALS

- ☒ RBC Lysis Buffer **BioLegend Catalog #420301**
- ☒ Anti-mouse CD3ε, clone 145-2C11 (LEAF™ format) **BioLegend Catalog #100314**
- ☒ Anti-mouse CD28, clone 37.51, (LEAF™ format) **BioLegend Catalog #102112**
- ☒ LEAF™ Purified anti-mouse IL-4 Antibody, clone 11B11 **BioLegend Catalog #504108**
- ☒ Recombinant Mouse IL-2 (carrier-free) **BioLegend Catalog #575402**
- ☒ Recombinant Mouse IL-12 (p70) (carrier-free) **BioLegend Catalog #577002**
- ☒ Monensin Solution (1,000X) **BioLegend Catalog #420701**
- ☒ Phorbol 12-myristate 13-acetate (PMA) **Sigma Aldrich Catalog #P8139**
- ☒ Ionomycin calcium salt from Streptomyces conglobatus **Sigma Aldrich Catalog #I0634**

Isolation of CD4+ Cells From Lymph Nodes

- 1 Harvest lymph nodes (superficial cervical, mandibular, axillary, inguinal, and mesenteric) from mice.

- 2 Tease lymph nodes through a sterile 70-µm nylon cell strainer to obtain single-cell suspensions incomplete RPMI containing 10% FCS (complete medium).
- 3 Resuspend cells in complete medium and use your favorite method to isolate CD4⁺ cells. Consider using our Mojosort™ Mouse CD4 T Cell Isolation Kit.

Th1 Polarization of CD4⁺ Cells

- 4 On day 0, coat 12-well plate with anti-mouse CD3ε, clone 145-2C11 (3µg/ml). Incubate at 37°C for 2 hours or 4°C overnight. Aseptically decant antibody solution from the plate. Wash plate 3 times with sterile PBS. Discard liquid.
02:00:00
- 5 Plate CD4⁺ cells at 1.0×10^6 /1ml/well. Culture cells for 5 days at 37°C, 5% CO₂, in the presence of anti-mouse CD28, clone 37.51 (3 µg/mL), anti-mouse IL-4, clone 11B11 (10 µg/mL), recombinant mouse IL-2 (5 ng/mL), and recombinant mouse IL-12 (10 ng/ml).
- 6 On day 3, if media is yellow, add 2 ml/well of fresh media.
- 7 On day 5, wash cells once and then restimulate in complete media with 50 ng/ml PMA, 1 µg/ml ionomycin and 10 µl monensin (1000x), in a 6-well plate in incubator at 37°C for 5 hours.
05:00:00
- 8 After harvesting, the cells are ready for staining.