



Th17 Polarization of Mouse CD4+ Cells V.3

DOI

dx.doi.org/10.17504/protocols.io.79vhr66



Sam Li¹

¹BioLegend

BioLegend

Tech. support email: tech@biolegend.com



Sam Li

BioLegend



DOI: dx.doi.org/10.17504/protocols.io.79vhr66

External link: <https://www.biolegend.com/protocols/th17-polarization-of-mouse-cd4--cells/4284/>

Protocol Citation: Sam Li . Th17 Polarization of Mouse CD4+ Cells. **protocols.io**

<https://dx.doi.org/10.17504/protocols.io.79vhr66>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Created: October 15, 2019

Last Modified: October 15, 2019

Protocol Integer ID: 28693

Keywords: th17, polarization, t cell



Materials

MATERIALS

✕ MojoSort™ Mouse CD4 T Cell Isolation Kit **BioLegend Catalog #480005, 480006, 480033**

✕ RBC Lysis Buffer **BioLegend Catalog #420301**

✕ Anti-mouse CD3ε clone 145-2C11 (Ultra-LEAF™ format) **BioLegend Catalog #100339**

✕ Anti-mouse CD28 clone 37.51 (Ultra-LEAF™ format) **BioLegend Catalog #102116**

✕ Anti-mouse IFN-γ clone XMG1.2 (Ultra-LEAF™ format) **BioLegend Catalog #505834**

✕ Anti-mouse IL-4 clone 11B11 (Ultra-LEAF™ format) **BioLegend Catalog #504122**

✕ Recombinant mouse IL-6 (carrier-free) **BioLegend Catalog #575704**

✕ Recombinant mouse IL-23 (carrier-free) **BioLegend Catalog #589002**

✕ Recombinant human TGF-β1 (carrier-free) **BioLegend Catalog #580702**

✕ Brefeldin A **BioLegend Catalog #420601**

✕ Monensin Solution **BioLegend Catalog #420701**



- Sterile PBS
- Cell culture medium (IMDM supplemented with 10% FBS)
- Sterile plastic petri dishes
- PMA (Phorbol 12-myristate 13-acetate) (Cat. No. P8139 from Sigma)
- Ionomycin (Cat. No. I0634 from Sigma)



Isolation of CD4⁺ Cells From Lymph Nodes

- 1 Harvest lymph nodes (superficial cervical, mandibular, axillary, inguinal, and mesenteric) from mice.
- 2 Tease lymph nodes through a sterile 70- μ m nylon cell strainer to obtain single-cell suspensions in complete IMDM containing 10% FCS (complete medium).
- 3 Resuspend cells in complete medium and use your favorite method to isolate CD4⁺ cells. Consider using our [**MojoSort™ Mouse CD4 T Cell Isolation Kit**](#).

Th17 Polarization of CD4⁺ Cells

- 4 On day 0, coat 60 × 15mm of plastic petri dishes with anti-mouse CD3 ϵ , clone 145-2C11 (5 μ g/ml). Incubate at 37°C for 2 hours or 4°C overnight. Aseptically decant antibody solution from the plate. Wash plate 3 times with sterile PBS. Discard liquid.
 02:00:00
- 5 Plate CD4⁺ cells at 10 × 10⁶/5 ml/dish. Culture cells for 4 days in the presence of anti-mouse CD28, clone 37.51 (5 μ g/mL), recombinant mouse IL-6 (50ng/mL), recombinant human TGF- β 1 (1ng/mL), recombinant mouse IL-23 (5ng/ml), anti-mouse IL-4 (10 μ g/mL), and anti-mouse IFN- γ (10 μ g/mL).
- 6 On day 3, slowly add 5ml of fresh media along with same the concentration of antibodies/cytokines as used on day 0.
- 7 On day 4, wash cells once and then restimulate in complete medium with 500ng/ml PMA and 500ng/mL ionomycin, in the presence of Brefeldin A (If you are looking for IL-21 production, use monensin) for 4-5 hours.
 05:00:00
- 8 After harvesting, the cells are ready for staining.

Tip: Recombinant human TGF- β is effective for stimulating mouse cells.