FEB 27, 2019

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.x8ufrww

Protocol Citation: Celeste M M. Karch, Rita Martinez, Jacob Marsh 2019. Thawing iPSC Plate. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.x8ufrww>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: Feb 17, 2019

Last Modified: Feb 27, 2019

PROTOCOL integer ID:
 20468

Thawing iPSC Plate

In 1 collection

Celeste M M. Karch¹, Rita Martinez¹, Jacob Marsh¹

¹Washington University in St Louis

Neurodegeneration Method Development Community

Tech. support email: ndcn-help@chanzuckerberg.com

Celeste M M. Karch

Washington University in St Louis

ATTACHMENTS

Comprehensive Genomic
 Editing and Screening
 Protocol Updated
 02142019.docx

GUIDELINES

This protocols is part of the [Screening Edited iPSC Clones collection](#).

SAFETY WARNINGS

Please refer to the SDS (Safety Data Sheet) for information about hazards, and to obtain advice on safety precautions.

- 1 Determine which clones to expand based on screening.

2 Coat 24 well plate with 250-300 ul Matrigel per well (1 well per well, from 96 well).

3 Incubate at 37 °C for 01:00:00 .

4 Remove Styrofoam box from -80 °C and remove plate.

Note

Check on plate after 00:15:00 to avoid over-thawing.

5 To each well add mTesR1 supplemented with 10 uM Rock inhibitor.

a. Remove desired cells from appropriate wells and transfer to 1.7 mL tube. Spin off freezing media. Plate cells in 24 well plate.

-or-

b. Remove desired cells from appropriate wells and dilute 1:5 (final volume of 500uL) in mTesR1 and plate in 24 well plate.

6 Incubate at 37 °C overnight.

7 Change mTesR1 daily.