FEB 27, 2019

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.x9afr2e

Protocol Citation: Celeste M M. Karch, Rita Martinez, Jacob Marsh 2019. Thawing Rosettes and NPC.
protocols.io
<https://dx.doi.org/10.17504/protocols.io.x9afr2e>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: Feb 17, 2019

Last Modified: Feb 27, 2019

PROTOCOL integer ID:
 20482

Thawing Rosettes and NPC

In 1 collection

Celeste M M. Karch¹, Rita Martinez¹, Jacob Marsh¹

¹Washington University in St Louis

Neurodegeneration Method Development Community

Tech. support email: ndcn-help@chanzuckerberg.com

Celeste M M. Karch

Washington University in St Louis

ATTACHMENTS

[IPSC CORTICAL DIFFERENTIATION 022017.pdf](#)

GUIDELINES

This protocol is part of the [IPSC CORTICAL DIFFERENTIATION](#) collection.

This method should be performed using sterile technique.

MATERIALS

Please refer to the attached full manuscript for required materials.

SAFETY WARNINGS

Please refer to the SDS (Safety Data Sheet) for information about hazards, and to obtain advice on safety precautions.

1 Aliquot 9 mL of pre-warmed DMEM/F12 into a 15 ml conical tube.

- 2 Remove cells from liquid nitrogen and thaw in 37 °C water bath for approximately 00:00:30 or until a small ice clump remains.
- 3 Add freshly thawed cells into the 9 mL of pre-warmed DMEM/F12 and mix gently by gently shaking tube 3 times. Avoid breaking up clumps of cells.
- 4 Centrifuge cells at 750 rpm for 00:03:00 . Aspirate supernatant.
- 5 Resuspend in 2 mL of NIM and plate into one well of a 6-well plate.

Note

If cell viability count is necessary, after resuspension remove 10 µl into a separate microfuge tube for cell viability count .