

Sep 05, 2020

The combination of LMWH (Low-Molecular-Weight Heparin) and Aspirin as a prophylaxis treatment for pre-eclampsia and SGA (Small-for-Gestational Age) neonates in medium- and high-risk women: protocol for a Systematic Review and Meta-Analysis.

DOI

dx.doi.org/10.17504/protocols.io.bhdtj26n

Christos-Georgios Kontovazainitis¹, Georgios Mitsiakos², Dimitra Gialamprinou²

¹Medical School, Aristotle University of Thessaloniki, University Campus, 54124 Thessaloniki, Greece; ²2nd Neonatal Department and Neonatal Intensive Care Unit (NICU), Aristotle University of Thessaloniki, Papageorgiou Hospital, Ring Road Nea Efkarpia, 56403, Thessaloniki, Greece



Christos-Georgios Kontovazainitis

OPEN ACCESS



DOI: dx.doi.org/10.17504/protocols.io.bhdtj26n

Document Citation: Christos-Georgios Kontovazainitis, Georgios Mitsiakos, Dimitra Gialamprinou 2020. The combination of LMWH (Low-Molecular-Weight Heparin) and Aspirin as a prophylaxis treatment for pre-eclampsia and SGA (Small-for-Gestational Age) neonates in medium- and high-risk women: protocol for a Systematic Review and Meta-Analysis.. **protocols.io** https://dx.doi.org/10.17504/protocols.io.bhdtj26n

License: This is an open access document distributed under the terms of the **Creative Commons Attribution License**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Created: June 10, 2020

Last Modified: September 05, 2020

Document Integer ID: 38035





The combination of LMWH (Low M...

DOWLOAD THE ATTACHED FILE (PDF) TO VIEW THE COMPLETE PROTOCOL.

Citation: Christos - Georgios Kontovazainitis, Georgios Mitsiakos, Dimitra Gialamprinou. The combination of LMWH (Low-Molecular-Weight Heparin) and Aspirin as a prophylaxis treatment for pre-eclampsia and SGA (Small-for-Gestational Age) neonates in medium- and high-risk women: protocol for a Systematic Review and Meta-Analysis.

Registration: Our protocol was registered with the International Prospective Register of Systematic Reviews (PROSPERO).

PROSPERO Registration number: CRD42020148500

This protocol is drafted based on the updated Preferred Reporting Items in Systematic Reviews and Meta Analyses (PRISMA) quidelines(1).

Keywords: pregnancy; high-risk pregnancy; pre-eclampsia; preeclampsia; aspirin; acetylsalicylic acid; 2acetoxybenzoic acid; heparin; low-molecular-weight heparin; LMWH; intrauterine growth restriction; IUGR; fetal growth restriction; small for gestational age; SGA; neonates; adverse events; perinatal mortality; systematic review; meta-analysis

Authors: Dr. Georgios Mitsiakos (GM), MD, PhD1

Dr. Christos-Georgios Kontovazainitis (C-GK), MSC²

Dr. Dimitra Gialamprinou (DG)¹

Affiliations:

¹2nd Neonatal Department and Neonatal Intensive Care Unit (NICU), Aristotle University of Thessaloniki, Papageorgiou Hospital, Ring Road Nea Efkarpia, 56403, Thessaloniki, Greece

²Medical School, Aristotle University of Thessaloniki, University Campus, 54124 Thessaloniki, Greece

Email:Dr. Georgios Mitsiakos MD, PhD: mitsiakos@auth.gr

Dr. Christos-Georgios Kontovazainitis, MSC: ckontova@auth.gr

Dr. Dimitra Gialamprinou: dgialamp@auth.gr

Corresponding author: Dr. Christos-Georgios Kontovazainitis 2nd Neonatal Department and Neonatal Intensive Care Unit Aristotle University of Thessaloniki Papageorgiou Hospital, Ring Road Nea Efkarpia

56403, Thessaloniki, Greece Tel: 0030 694 538 3092

Email: ckontova@auth.gr



Author contributions: C-GK, GM and DG drafted the manuscript. C-GK, GM and DG contributed to the development of the selection criteria, the risk of bias assessment strategy and data extraction criteria. C-GK developed the search strategy. GM provided expertise on neonatology field. C-GK provided expertise on the field of obstetrics-gynaecology. All authors read, provided feedback and approved the final manuscript.

Amendments: In the event of amendments in the protocol, the date of each amendment will be accompanied by an explanation of the alterations and the relevant rationale.

Version 1.0 12.02.2020

Version 2.0 10.06.2020

- ·Changes in the inclusion criteria: studies that only include the secondary outcomes and not the primary ones will not be excluded. Case-control studies will be excluded.
- ·Changes concerning the tools used for the assessment of the risk of bias.
- ·Appendix 4 added.

Financial support, funding sources: The authors declare no source of institutional or external financial support in relation to this work.