VERSION 4

DEC 14, 2018

OPEN ACCESS

DOI:

[dx.doi.org/10.17504/protocols.io.wdtfa6n](https://doi.org/10.17504/protocols.io.wdtfa6n)

External link:

<https://doi.org/10.1371/journal.pone.0217633>

Protocol Citation: Quentin M. Nunes, Dunhao Su, Phuong Thao Bui, Deborah M. Simpson, Philip J. Brownridge, Changye Sun, Yong Li, Zhang X, Wei Huang, Daniel J. Rigden, Robert J. Beynon, Robert Sutton, David G. Fernig 2018. The heparin-binding proteome in normal pancreas and murine experimental acute pancreatitis. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.wdtfa6n>

MANUSCRIPT CITATION:

Nunes QM, Su D, Brownridge PJ, Simpson DM, Sun C, Li Y, Bui TP, Zhang X, Huang W, Rigden DJ, Beynon RJ, Sutton R, Fernig DG (2019) The heparin-binding proteome in normal pancreas and murine experimental acute pancreatitis. PLoS ONE 14(6): e0217633. doi: [10.1371/journal.pone.0217633](https://doi.org/10.1371/journal.pone.0217633)

The heparin-binding proteome in normal pancreas and murine experimental acute pancreatitis V.4

PLOS One

Quentin M. Nunes¹, Dunhao Su², Phuong Thao Bui², Deborah M. Simpson³, Philip J. Brownridge³, Changye Sun⁴, Yong Li^{5,6,7}, Zhang X¹, Wei Huang⁸, Daniel J. Rigden², Robert J. Beynon³, Robert Sutton¹, David G. Fernig²

¹Liverpool Pancreatitis Research Group, Department of Molecular and Clinical Cancer Medicine, University of Liverpool;

²Department of Biochemistry, Institute of Integrative Biology, University of Liverpool;

³Protein Function Group, Institute of Integrative Biology, University of Liverpool & Department of Biochemistry, Institute of Integrative Biology, University of Liverpool;

⁴Department of Biochemistry, Institute of Integrative Biology, University of Liverpool & Henan Key Laboratory of Medical Tissue Regeneration, Xinxiang Medical University, Xinxiang, Henan, 453003, China;

⁵Department of Biochemistry, Institute of Integrative Biology, University of Liverpool & College of Life and Environmental Science;

⁶Wen Zhou University; ⁷Wenzhou, China;

⁸Liverpool Pancreatitis Research Group, Department of Molecular and Clinical Cancer Medicine, University of Liverpool & Department of Integrated Traditional Chinese and Western Medicine, Sichuan Provincial Pancreatitis Centre and West China-Liverpool Biomedical Research Centre, West China Hospital, Sichuan University, Chengdu, China

Phuong Thao Bui

University of Liverpool

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Dec 12, 2018

Last Modified: Dec 14, 2018

PROTOCOL integer ID:
18579

ABSTRACT

Acute pancreatitis (AP) is acute inflammation of the pancreas, mainly caused by gallstones and alcohol, driven by changes in communication between cells. Heparin-binding proteins (HBPs) play a central role in cell communication. Therefore, we used heparin affinity proteomics to identify extracellular HBPs in pancreas and plasma of normal mice and in a caerulein mouse model of AP. Many new extracellular HBPs (360) were discovered in the pancreas, taking the total number of HBPs known to 786. Extracellular pancreas HBPs form highly interconnected protein-protein interaction networks in both normal pancreas (NP) and AP. Thus, HBPs represent an important set of extracellular proteins with significant regulatory potential in the pancreas. HBPs in NP are associated with biological functions such as molecular transport and cellular movement that underlie pancreatic homeostasis. However, in AP HBPs are associated with additional processes such as acute phase response signalling, complement activation and mitochondrial dysfunction. Plasma HBPs in AP included known AP biomarkers such as serum amyloid A, as well as emerging targets such as histone H2A. Pancreas HBPs are extracellular and so easily accessible and are potential drug targets in AP, whereas plasma HBPs represent potential biomarkers for AP. These need further investigation for potential applications in the management of AP.

MATERIALS

STEP MATERIALS

✕ RapiGest SF Surfactant **Contributed by users Catalog #23221**

✕ Iodoacetamide **P212121 Catalog #AK-U470**

✕ DTT **Merck MilliporeSigma (Sigma-Aldrich) Catalog #D0632**

✕ Trypsin from porcine pancreas **Merck MilliporeSigma (Sigma-Aldrich) Catalog #T7409**

✕ Trifluoroacetic acid (TFA) **Contributed by users**

Blood obtained from 6-8 week old male adult CD1 mice (weight range, 24-30 g) with normal pancreas (NP) or experimental acute pancreatitis (AP), induced as described above, was collected in 0.38% (w/v) sodium citrate (Sigma-Aldrich, Gillingham, UK) and centrifuged at 5,000 g at 4 °C for 15 min [27]. Supernatant plasma was extracted and frozen at -20 °C.

Heparin affinity chromatography

- 1 The frozen murine plasma was defrosted on ice and centrifuged at 16,100 g for 10 min, at 4 °C.

🌡 4 °C 16,100 g ⌚ 00:10:00

The resulting supernatant was diluted (1:8) in 75 mM NaCl, 6.85 mM Na₂HPO₄, 3.15 mM

NaH₂PO₄, pH 7.2.

Note

Note

This method was effective in reducing the high abundance protein, albumin, down to levels that would not affect the MS analysis of plasma HBPs.

The diluted supernatant was centrifuged at 5,000 g for 5 min.

00:05:00 5000 g

and the resulting supernatant was then applied on a 1 mL Hi-Trap heparin column

Equipment

1 mL Hi-Trap heparin column

NAME

Affinity column

TYPE

GE Healthcare Life Sciences

BRAND

17040601

SKU

<https://www.gelifesciences.com/en/br/shop/chromatography/resins/affinity-specific-groups/hitrap-heparin-hp-affinity-columns-p-01737>

LINK

equilibrated in 75 mM NaCl, 6.85 mM Na₂HPO₄, 3.15 mM NaH₂PO₄, pH 7.2.

The heparin column was subsequently washed extensively with 50 mL of **modified PBS** (150

mM NaCl, 13.7 mM Na₂HPO₄, 6.3 mM NaH₂PO₄, pH7.2). The heparin-bound fraction was then eluted with 2 M NaCl in **modified PBS**.

Note

The receipt for PBS has been modified here. Please check carefully.

[M] 2 Molarity (M) NaCl

Protein adsorption using StrataClean for mass spectrometry

- 2 In order to eliminate the high concentration of electrolytes (2.15 M NaCl) in the eluate, so as not to influence MS analysis, StrataClean™ beads (Agilent Technologies, UK) were used to adsorb proteins.

⊗ StrataClean™ beads **Contributed by users Catalog #400714**

Briefly, 100 µg of the heparin-bound fraction was mixed with 30 µL StrataClean slurry, vortexed for 2 min. and centrifuged at 2,000 g for 2 min at 4 °C.

🌡 4 °C ⌚ 00:02:00

The supernatant was removed carefully and discarded.

The pellet was first washed with PBS and then further washed twice with H₂O.

Sample preparation for mass spectrometry

- 3 In the case of plasma samples, the StrataClean™ beads were resuspended in 80 µL of 25 mM ammonium bicarbonate and 5 µL of 1 % (w/v) Rapigest (Waters, Manchester, UK) added and the samples shaken at 450 rpm for 10 min at 80 °C.

RapiGest SF Surfactant **Contributed by users** Catalog #23221

80 °C for the denature by RapiGest 00:10:00

Samples were reduced by the addition of 5 µL of 60 mM (DTT) and incubated at 60°C 60 °C for 10 min 00:10:00 and alkylated (addition of 5 µL of 180 mM iodoacetamide and incubation at room temperature for 30 min in the dark).

Iodoacetamide P212121 Catalog #AK-U470

DTT Merck MilliporeSigma (Sigma-Aldrich) Catalog #D0632

Trypsin (Sigma, Poole, UK, proteomics grade) was reconstituted in 50 mM acetic acid to a concentration of 0.2 µg/µL and 5 µL (1 µg) added to the sample followed by overnight incubation at 37 °C. 17:00:00 overnight 37 °C

Trypsin from porcine pancreas Merck MilliporeSigma (Sigma-Aldrich) Catalog #T7409

Note

Samples were mixed on a rotating mixer

The following day the digestion was terminated and Rapigest removed by acidification with TFA (1 µL)

Trifluoroacetic acid (TFA) **Contributed by users**

and incubation at 37 °C for 45 min. 37 °C 00:45:00

Samples were centrifuged at 17,200 g for 30 min and the clarified supernatants transferred to tubes.

00:30:00 17,200 g centrifuge