


Apr 14, 2020

The role of multiprotein bridging factor 1 (MBF1c) in Funariaceae species under salt stress

DOI

dx.doi.org/10.17504/protocols.io.be3sjgne



Maria Victória Magalhães de Vargas^{1,2}, Guilherme Afonso Kessler de Andrade^{1,2}, Sara Navarrete Bohi Goulart^{1,2}, Bruna Mota Bernardes^{1,2}, Filipe Victoria^{1,2}

¹Núcleo de Estudos da Vegetação Antártica (NEVA), Universidade Federal do Pampa (UNIPAMPA), São Gabriel, RS, Brazil;

²neva@unipampa.edu.br



Filipe Victoria

OPEN  ACCESS



DOI: dx.doi.org/10.17504/protocols.io.be3sjgne

Protocol Citation: Maria Victória Magalhães de Vargas, Guilherme Afonso Kessler de Andrade, Sara Navarrete Bohi Goulart, Bruna Mota Bernardes, Filipe Victoria 2020. The role of multiprotein bridging factor 1 (MBF1c) in Funariaceae species under salt stress. [protocols.io https://dx.doi.org/10.17504/protocols.io.be3sjgne](https://dx.doi.org/10.17504/protocols.io.be3sjgne)

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Other

We attempted this protocol, but could not get it to work in our group

Created: April 14, 2020

Last Modified: April 14, 2020

Protocol Integer ID: 35666

Keywords: Abiotic Stress, expression factors, *Funaria hygrometrica*, *Physcomitrella patens*, *Physcomitrium acutifolium*, qPCR,

Abstract

Bryophytes are one of the most successful plants in nature and can tolerate several environmental stresses, such as saline stress. MBF1 is a transcription factor that may be involved in reprogramming transcription, which in turn may confer tolerance to stress conditions. The objective of this study was to quantify PaMBF1c under saline stress in three species of moss: *Funaria hygrometrica*, *Physcomitrium acutifolium*, and *Physcomitrella patens*. All moss species underwent 12 h and 24 h treatment with salt at different concentrations, followed by qRT-PCR experiments.

In *F. hygrometrica*, the MBF1c gene was up-regulated when subjected to 12 h salt treatment at a medium concentration, and may be an important factor for the tolerance of plants to NaCl concentrations. In *P. acutifolium*, 24 h NaCl treatment at of 400, 600, and 800 mM resulted in a gradual up-regulation of MBF1c expression, reflective of the increasing NaCl concentration. In *Physcomitrella*, the expression profile of MBF1c showed a down-regulation when treated with 200, 400, and 800 mM NaCl, suggesting that MBF1c may not interfere with the saline stress response in this plant.

The diversity of these expression profiles demonstrate that the MBF1c response is not universal neither among plants nor between species of the same family, as shown in the mosses studied.

Attachments



[Methods_Protocol.doc...](#)

14KB

Materials

MATERIALS

-  Qubit™ dsDNA HS Assay Kit Invitrogen - Thermo Fisher Catalog #Q32851
-  GoTaq(R) Hot Start Polymerase, 2500u Promega Catalog #M5006
-  Qubit RNA HS Assay Kit Thermo Fisher Scientific Catalog #Q32852
-  PureLink™ RNA Mini Kit Thermo Fisher Scientific Catalog #12183020
-  High-Capacity RNA-to-cDNA™ Kit Thermo Fisher Catalog #4387406

