VERSION 1
MAY 16, 2020

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.7r6hm9e

External link:
<https://neb.com/protocols/2019/03/19/thermolabile-proteinase-k-typical-reaction-protocol>

Protocol Citation: New England Biolabs 2020. Thermolabile Proteinase K Typical Reaction Protocol. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.7r6hm9e>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Sep 27, 2019

Last Modified: May 16, 2020

Thermolabile Proteinase K Typical Reaction Protocol V.1

New England Biolabs¹

¹New England Biolabs

New England Biolabs (NEB)

Tech. support phone: +1(800)632-7799 email: info@neb.com

New England Biolabs

New England Biolabs

MATERIALS

MATERIALS

Thermolabile Proteinase K New England Biolabs Catalog #P8111S

SAFETY WARNINGS

Please see SDS (Safety Data Sheet) for hazards and safety warnings.

- 1 Reactions may be scaled-up linearly to accommodate larger amounts of substrate and larger reaction volumes. Optimal buffering reagents, enzyme quantity, incubation temperatures and times may vary for particular substrates. Typical restriction enzyme cleanup conditions are as follows:

To a 50 μL restriction enzyme digest containing 1 μg of DNA and 10 units of a restriction enzyme:

- 2 Add 1 μL of *Thermolabile Proteinase K*, mix gently.

- 3 Incubate reaction at 37 °C for 00:15:00 .

- 4 Incubate reaction at 55 °C for 00:10:00 to inactivate *Thermolabile Proteinase K*.

Note

Note: Optimal reaction buffers, reaction temperatures as well as additional reaction properties of *Thermolabile Proteinase K* can be found in detail in the [FAQ](#). Optimization may be required for other cleanup applications (ligation, extension, PCR, etc.)