DEC 01, 2017

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.kpxcvpn

Protocol Citation: Tomaz Crochemore, Thiago Domingos Corrêa, Marcus D. Lancé, Cristina Solomon, Ary Serpa Neto, João Carlos de Campos Guerra, Priscila Scolmeister Lellis, Livia Muller Bernz, Natalia Nunes, Cassio Massashi Mancio, Ana Paula Hitomi Yokoyama, Eliézer Silva 2017.

THROMBOELASTOMETRY
 PROFILE IN CRITICALLY ILL
 PATIENTS. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.kpxcvpn>

MANUSCRIPT CITATION:

References

1. Palta S, Saroa R, Palta A. Overview of the coagulation system. Indian J Anaesth. 2014;58(5):515-23.
2. Jaffer U, Wade RG, Gourlay T. Cytokines in the systemic inflammatory response syndrome: a review. HSR Proc Intensive Care Cardiovasc Anesth. 2010;2(3):161-75.
3. Lier H, Vorweg M, Hanke A, Görlinger K. Thromboelastometry guided therapy of severe bleeding. Essener Runde Hamostaseologie. 2013;33(1):51-61.
4. Ng VL. Liver disease,

THROMBOELASTOMETRY PROFILE IN CRITICALLY ILL PATIENTS

Tomaz Crochemore¹, Thiago Domingos Corrêa¹, Marcus D. Lancé², Cristina Solomon³, Ary Serpa Neto¹, João Carlos de Campos Guerra⁴, Priscila Scolmeister Lellis⁴, Livia Muller Bernz¹, Natalia Nunes¹, Cassio Massashi Mancio⁵, Ana Paula Hitomi Yokoyama⁶, Eliézer Silva¹

¹1. Hospital Israelita Albert Einstein, Intensive Care Unit, São Paulo, Brazil;

²2. Anaesthetics, Cardiothoracic Surgery, Emergency Medicine. Maastricht University Medical Centre, Maastricht, The Netherlands;

³3. Paracelsus Medical University, Salzburg, Austria, and Octapharma AG, Lachen, Switzerland;

⁴4. Hospital Israelita Albert Einstein, Laboratory Center, São Paulo, Brazil;

⁵5. Hospital Israelita Albert Einstein, Pharmacy, São Paulo, Brazil;

⁶6. Hospital Israelita Albert Einstein, Blood Bank, São Paulo, Brazil

Tomaz Crochemore

coagulation testing, and hemostasis. Clin Lab Med. 2009;29(2):265-82.

5. Levi M, Schultz M, van der Poll T. Coagulation biomarkers in critically ill patients. Crit Care Clin. 2011;27(2):281-97.
6. Meybohm P, Zacharowski K, Weber CF. Point-of-care coagulation management in intensive care medicine. Crit Care. 2013;17(2):218.
7. Benes J, Zatloukal J, Kletecka J. Viscoelastic Methods of Blood Clotting Assessment - A Multidisciplinary Review. Front Med (Lausanne). 2015;2:62.
8. Sarani B, Dunkman WJ, Dean L, Sonnad S, Rohrbach JI, Gracias VH. Transfusion of fresh frozen plasma in critically ill surgical patients is associated with an increased risk of infection. Crit Care Med. 2008;36(4):1114-8.
9. Spiess BD, Royston D, Levy JH, Fitch J, Dietrich W, Body S, et al. Platelet transfusions during coronary artery bypass graft surgery are associated with serious adverse outcomes. Transfusion. 2004;44(8):1143-8.
10. Murphy GJ, Reeves BC, Rogers CA, Rizvi SI, Culliford L, Angelini GD. Increased mortality, postoperative morbidity, and cost after red blood cell transfusion in patients having cardiac surgery. Circulation. 2007;116(22):2544-52.
11. Haas T, Görlinger K, Grassetto A, Agostini V, Simioni P, Nardi G, et al. Thromboelastometry for guiding bleeding management of the critically ill patient: a systematic review of the literature. Minerva Anestesiol. 2014;80(12):1320-35.
12. Weber CF, Görlinger K, Meininger D, Herrmann E, Bingold T, Moritz A, et al. Point-of-care testing: a prospect

ABSTRACT

ABSTRACT

BACKGROUND: Transfusion therapy is associated with increased morbidity, mortality and costs. Conventional coagulation tests are weak bleeding predictors, poorly reflecting coagulation in vivo. Thromboelastometry (ROTEM) provides early identification of coagulation disorders and can guide transfusion therapy by goals, reducing blood components transfusion. **Objective:** The aim of this study is to describe coagulation profile of critically ill patients using ROTEM and evaluate the association between CCT and thromboelastometry. **METHODS:** This is a retrospective, observational study conducted in medical-surgical intensive care unit (ICU). Adult patients (≥ 18 years) admitted to ICU between November 2012 and December 2014, in whom ROTEM analyses were performed for diagnosis of coagulopathy, were included in this study. The first ROTEM and CCT after ICU admission were recorded simultaneously. Additionally, we collected data on blood components transfusion and hemostatic agents immediately after laboratory tests results. **RESULTS:** The study included 531 patients. Most ROTEM tests showed normocoagulability profile [INTEM (54.8%), EXTEM (54.1%) and FIBTEM (53.3%)] with divergent results in relation to CCT: low platelet count (51.8% in INTEM and 55.9% in EXTEM); prolonged aPTT (69.9% in INTEM and 63.7% in EXTEM) and higher INR (23.8% in INTEM and 27.4% in EXTEM). Approximately 16,7% of patients with normocoagulability in ROTEM received platelet concentrates and 10% fresh frozen plasma. **CONCLUSION:** The predominant ROTEM profile observed in this sample of critically ill patients was normocoagulability in contrast to CCT, suggesting coagulopathy and leading to a possibly unnecessary allogenic blood component transfusion. ROTEM test may avoid inappropriate allogeneic blood products transfusion in these patients.

ATTACHMENTS

[data protocol.xlsx](#)

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Nov 08, 2017

Last Modified: Mar 01, 2018

PROTOCOL integer ID:
8663

Keywords:
thrombelastometry,
thromboelastography,
hemostatic therapy,
hematologic tests, blood
coagulation tests, massive
transfusion, blood product