

Mar 06, 2020

## Total Phenolic Content (TPC)

 [PLOS One](#)

DOI

[dx.doi.org/10.17504/protocols.io.bda5i2g6](https://dx.doi.org/10.17504/protocols.io.bda5i2g6)

Jing Xu<sup>1</sup>

<sup>1</sup>Hainan University



Jing Xu

Hainan University

---

OPEN  ACCESS



DOI: [dx.doi.org/10.17504/protocols.io.bda5i2g6](https://dx.doi.org/10.17504/protocols.io.bda5i2g6)

External link: <https://doi.org/10.1371/journal.pone.0234435>

**Protocol Citation:** Jing Xu 2020. Total Phenolic Content (TPC). **protocols.io**  
<https://dx.doi.org/10.17504/protocols.io.bda5i2g6>

**Manuscript citation:**

Zhou J, Yang Q, Zhu X, Lin T, Hao D, Xu J (2020) Antioxidant activities of *Clerodendrum cyrtophyllum* Turcz leaf extracts and their major components. PLoS ONE 15(6): e0234435. doi: [10.1371/journal.pone.0234435](https://doi.org/10.1371/journal.pone.0234435)

**License:** This is an open access protocol distributed under the terms of the **[Creative Commons Attribution License](#)**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

**Protocol status:** Working

**We use this protocol and it's working**

**Created:** March 06, 2020

**Last Modified:** March 06, 2020

**Protocol Integer ID:** 33853



## Abstract

The TPC from samples was determined by a colorimetric method based on the procedure described by Zhou et al. [10] 2 ml Folin-Ciocalteu (FC) reagent was added to 2 ml of diluted extracts. After 3 min, 750  $\mu$ L of sodium carbonate anhydrous solution (7.5%, w/v) was added, and the mixture was adjusted to 10 ml with distilled water. After 2 h, the absorbance was recorded at 765 nm. Calibration curves were constructed with gallic acid as a standard at concentrations ranging from 0–100 mg/ml. ml.

