OCT 22, 2019

Total RNA Isolation from Bulk Tissue or Isolated Cells

Brad Godfrey¹

¹University of Michigan - Ann Arbor

KPMP

Tech. support email: info@kpmp.org

Brad Godfrey

OPEN ACCESS

DOI:

dx.doi.org/10.17504/protocols.io.8dehs3e

External link:

<http://www.kpmp.org>

Protocol Citation: Brad Godfrey 2019. Total RNA Isolation from Bulk Tissue or Isolated Cells. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.8dehs3e>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Oct 17, 2019

Last Modified: May 05, 2020

PROTOCOL integer ID:
28806

Procedures

- 1 Transfer biopsy in 1.8 mL cryotube (polypropylene) in 1.0 mL RNAlater
- 2 Use color coding cap inserts to indicate biopsy origin
- 3 Store at -20 °C until RNA extraction
- 4 Perform microdissection of glomeruli / tubulointerstitium
- 5 Store dissected tissue portion in less than 50 µL RNAlater solution and store at -20 °C until total RNA extraction
- 6 For total RNA/DNA extraction use “AllPrep DNA/RNA Micro Kit” (# 80284) from Qiagen following respective protocol. Use optional Beta-ME for glomerular compartment.

Note: Elute RNA in 16 µL H₂O in nucleotide low-binding tubes (Eppendorf #022431021) and store at -80 °C .

Note: Keep first flow-through (< 80 bases) for future analysis of small RNAs.

Note

Quality and quantity were measured using Agilent Bioanalyzer 2100

Elute RNA in 16µL

cDNA Generation

- 7 Use NEBNext® Ultra™ II Directional RNA Library Prep Kit for Illumina® (rRNA depletion workflow)1. Use NEBNext® Ultra™ II Directional RNA Library Prep Kit for Illumina® (rRNA depletion workflow)
- 8 1.Store cDNA at -20 °C

NGS Library Preparation

- 9 Use NovaSeq 6000 to sequence. (The University of Michigan Advanced Genomics Core)1. Use NovaSeq 6000 to sequence. (The University of Michigan Advanced Genomics Core)