JUL 10, 2019

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.457gy9n

Protocol Citation: Antonio Mondini, Morten Dencker Schostag, Lea Ellegaard-Jensen, Toke Bang-Andreasen, Muhammad Zohaib Anwar, Cristina Purcarea, Carsten Suhr Jacobsen 2019. Total RNA Protocol (extraction, quantification and Illumina library preparation).

protocols.io
<https://dx.doi.org/10.17504/protocols.io.457gy9n>

🌐 Total RNA Protocol (extraction, quantification and Illumina library preparation)

Antonio Mondini¹, Morten Dencker Schostag²,
 Lea Ellegaard-Jensen³, Toke Bang-Andreasen³,
 Muhammad Zohaib Anwar³, Cristina Purcarea¹,
 Carsten Suhr Jacobsen³

¹Laboratory of Molecular Microbiology, Institute of Biology Bucharest, Department of Microbiology, Romanian Academy, Bucharest, Romania;

²Geochemical Department, Geological Survey of Denmark and Greenland, Øster Voldgade 10, 1350 Copenhagen, Denmark;

³Department of Environmental Sciences, Aarhus University, 399 Frederiksborgvej, 4000 Roskilde, Denmark

Muhammad Zohaib Anwar
 Aarhus University

ABSTRACT

This experimental procedure was designed to isolate total the RNA aiming to obtain intact and high-quality integrity (RIN>7) for further next generation sequencing analysis. The protocol was developed to analyse the environmental microbial active community providing a clear overview of the microbial transcriptional activity in a specific time or after a different condition exposition sets, such as the investigation of thermal adaptation after heat-shock. The procedure is specifically designed to extract total-RNA from water samples although it can be used for various environmental samples as done in Schostag et al. and Bang-Andreasen et al. 2019 in soil with similar accuracy results. The total-RNA yield was >1000 ng/mL with a RIN ranging from 7 to 8.5. The final Illumina library preparation yield ranged between 15 and 31 ng/μL.

This protocol is a compiled by using the Qiagen RNeasy® PowerSoil® Total RNA Kit with additional steps according the DNeasy Max®Kit (Qiagen), NEBNext® Ultra II Directional RNA Library Prep Kit for Illumina® (New England Biolabs, Inc), KAPA Library Quantification Kit for Illumina® Platforms (Kapa Biosystems), Qubit 2.0 (Thermo Fisher Scientific, Hvidovre, Denmark), Bioanalyzer® (Agilent Technologies, Inc.) sample Purification Beads (New England Biolabs) and Liquid G2 DNA/RNA Enhancer (Ampliqon) manufacturer protocols.

MANUSCRIPT CITATION:

1) Schostag M, Priemé A, Jacquioud S, Russel J, Ekelund F, Jacobsen CS. Bacterial and protozoan dynamics upon thawing and freezing of an active layer permafrost soil. The ISME journal. 2019 May;13(5):1345. 2) Bang-Andreasen T, Anwar MZ, Lanzén A, Kjølner R, Rønn R, Ekelund F and Jacobsen CS (2019) Total RNA-sequencing reveals multi-level microbial community changes and functional responses to wood ash application in agricultural and forest soil. DOI: <https://doi.org/10.1101/621557> 3) Schostag MD, Anwar MZ, Jacobsen CS, Larose C, Vogel TM, Maccario L, Jacquioud S, Faucherre S and PrieméA (2019) Transcriptomic responses to warming and cooling of an Arctic tundra soil microbiome. DOI: <https://doi.org/10.1101/599233>

3) Schostag MD, Anwar MZ, Jacobsen CS, Larose C, Vogel TM, Maccario L, Jacquioud S, Faucherre S and PrieméA (2019) Transcriptomic responses to warming and cooling of an Arctic tundra soil microbiome. DOI: <https://doi.org/10.1101/599233>

3) Schostag MD, Anwar MZ, Jacobsen CS, Larose C, Vogel TM, Maccario L, Jacquioud S, Faucherre S and PrieméA (2019) Transcriptomic responses to warming and cooling of an Arctic tundra soil microbiome. DOI: <https://doi.org/10.1101/599233>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol in our group at Department of Environmental Sciences, Aarhus University and other collaborators and it is working very well

Created: Jul 04, 2019

Last Modified: Jul 10, 2019

PROTOCOL integer ID:
25503

Keywords: Total-RNA, Environmental Microbiomes, Illumina, mRNA, rRNA

GUIDELINES

Perform all the work under a microbiological hood and change gloves frequently.

MATERIALS

Figure 1: Workflow Scheme

MATERIALS

- ☒ NEBNext Multiplex Oligos for Illumina (Index Primers Set 1) - 24 rxns **New England Biolabs Catalog #E7335S**
- ☒ Liquid nitrogen **Contributed by users**
- ☒ UltraPure™ Phenol:Chloroform:Isoamyl Alcohol (25:24:1, v/v) **Thermo Fisher Scientific Catalog #15593031**
- ☒ NEBNext Ultra II Directional RNA Library Prep Kit for Illumina - 24 rxns **New England Biolabs Catalog #E7760S**
- ☒ Disposable Latex Gloves, Medium, 100/Box **Bio Basic Inc. Catalog #GL002M.SIZE.1PK**
- ☒ Qubit dsDNA HS Assay kit **Thermo Fisher Scientific Catalog #Q32854**
- ☒ RNase Zap **Sigma Aldrich Catalog #R2020-250ML**
- ☒ Bioanalyzer chips and reagents (DNA High Sensitivity and small RNA kit) **Agilent Technologies Catalog #5067-1548**
- ☒ 80% Ethanol **Contributed by users**
- ☒ NEBNext Sample Purification Beads **New England Biolabs Catalog #E7767**
- ☒ Ice & ice bucket **Contributed by users**
- ☒ Qubit RNA HS Assay Kit **Thermo Fisher Scientific Catalog #Q32852**
- ☒ RNeasy® PowerSoil® Total RNA Kit **Qiagen Catalog #12866-25**
- ☒ S-Pak Filters 0.22µm 47mm white gridded **Merck Millipore Catalog #GSWG047S6**
- ☒ DNeasy Max®Kit **Qiagen Catalog #15200-50**
- ☒ KAPA Library Quantification Kit for Illumina® Platforms **Kapa Biosystems Catalog #KK4835**
- ☒ Liquid G2 DNA/RNA Enhancer **Ampliqon Catalog #A420015**

1. Microbiological Hood
2. Filtering system
3. Vacuum pump
4. Vortex-Genie® 2 Vortex
5. Vortex Adapter for 4 (15 ml) tubes (cat. no. 13000-V1-15)
6. Pipette tips with filter (20 µL – 1000 µL)
7. Thermoblock (set at 45°C)
8. Incubator for -20 and 4°C
9. 96-well 0.2 ml PCR Plates and Microseal® 'B' Adhesive Sealer (BioRad MSB-1001) or 0.2 ml RNase-free tube
10. 15 mL tubes and 1.5 mL vials racks
11. Qubit Invitrogen
12. Biology Tweezer 115mm, Ideal Tak
13. Bioanalyzer, Agilent G2991AA
14. SimpliAmp Thermal Cycler PCR, Applied Biosystems
15. Benchtop Centrifuge

16. Mini-centrifuge
17. Magnetic Stand

SAFETY WARNINGS

- ! When using Phenol Chloroform work under the fume hood, when not possible, seal the tubes properly with parafilm

BEFORE START INSTRUCTIONS

Remove RNases cleaning the surfaces, pipettes/forceps and gloves using RNaseZAP according to the manufacturer protocol. Pre-cool all the centrifuges.

Water samples filtration

- 1 Assemble the filtration system and place the 0.22 µm MF-membranes filters using RNase free forceps
- 2 Once completed the filtration, transfer the filters in the 15 mL tubes with beads (provided in the RNeasy@kit).

Note

Transfer the filters carefully folding it to fit in the 15mL tubes using two forceps and letting the surface with visible filtered particles inward the tube. Push carefully down to the beads.

Safety information

No more than 2 filters can be added to a single tube

RNA Isolation

- 3 Add G2 and snap freeze immersing the bead tubes 15 seconds in liquid nitrogen keeping the lid slightly open

Note

Add 500µL of G2 each filter. If a thick sediment layer accumulates on the filter, it has to be weighed and 500 µL of G2 added each 0.25 grams of sediments. No more than 1 gram can be added to the tube. Addition of more soil/sediments can reduce the lyse cells efficiency. For more info on G2 test see appendix 1 ➡ go to step #133 . Alternatively, to liquid nitrogen, tubes can be freeze O/N at -20°C

4 Immediately add 2.5 ml of PowerBead Solution

5 Add 0.25 ml of Solution SR1

6 Add 0.8 ml of Solution IRS

7 Add 3.5 ml of phenol/chloroform/isoamyl alcohol (pH 6.5) and vortex until the solution is homogeneous.

Note

After vortexing the solution should have a muddy consistency. At this stage the tubes have to be placed outside the Microbiological hood and to avoid phenol chloroform diffusion in the air is preferably to seal the tubes using parafilm

8 Place the PowerBead tube horizontally on a Vortex Adapter (cat. no. 13000-V1-15) and vortex at maximum speed for 20 minutes at 🌡 Room temperature

Note

This vortexing time was specifically adapted for tubes containing filters with the addition of Phenil chloroform and previously snap frozen in Liquid Nitrogen. For different samples vortex for 15 minutes according to the manufacturer protocol

9

Remove PowerBead tube from the shaker and centrifuge Room temperature for 10 minutes at 2,500 x g

Note

For different centrifuge unit see appendix 2 [go to step #134](#)

- 10 Transfer the upper aqueous phase (avoid the interphase and lower phenol layer) to a clean 15 ml Collection Tube (provided) using a 1000µL pipette. Ensure to take only the top layer (Approx. 3 mL) leaving some aqueous phase to not disturb the interphase.

Note

If the filters, after centrifuge, are interfering with normal procedure of pipetting or have formed agglomerates, pour the tube content (leaving behind the filters) in a new 15 mL tube and repeat the centrifugation at step 9.

- 11 Add 1.5 ml of Solution SR3 to the tube containing the aqueous phase and vortex to mix until the solution gets homogeneous

- 12 Incubate at 4 °C for 10 minutes

13 Centrifuge at 2,500 x g Room temperature for 10 minutes

14 Transfer the supernatant, without disturbing the pellet if present (usually not present), to a new 15 ml Collection Tube (provided)

Note

Its preferred to leave behind few microliters

15 Add 5 mL of Solution SR4 to the supernatant in the Collection Tube using a serological 5 mL pipette

16 Vortex until the biphasic layer disappear (2 cycles of 10 seconds)

17 Incubate at Room temperature for 30 minutes

18 Centrifuge at 2500 x g at Room temperature for 30 minutes

19 Decant the supernatant

20 Quickly spin down the tubes to collect remaining drops on the tubes wall

21 Remove the collected liquid without disturbing the pellet

22 Allow the tubes to dry for 5 minutes

Note

Do not let the pellet over dry (more than 10 minutes) or will result hard to resuspend it. Pellet colors were observed in a range of light grey and more often light/dark brown.

RNA Purification

23 Shake Solution SR5 and add 1 ml to the 15 ml Collection Tube.

24 Resuspend the pellet completely by repeatedly pipetting.

Note

Usually is hard to be resuspended and using a P1000 pipette just remove the pellet from the wall and proceed to the next step.

25 Incubate the tubes at **45 °C** in a preheated thermoblock. Check on the tube every 5 min and vortex the solution. Visual inspection should reveal the total resuspension.

26 Repeat step 25 until the pellet is fully resuspended

27 Once the pellet is resuspended place the JetStar Mini Column (provided capture column) in a new 15 ml tube

28 Shake SR5 solution and add 2 mL to the RNA capture column

Safety information

From this step on never allow to completely gravity flow through the column. A dry column will reduce the downstream RNA yield. Add warning here

29 Once the SR5 solution has reached the top of the resin (1 mm thickness) in the column add the dissolved pellet to the column and allow it to gravity flow through the column

30 When dissolved pellet has reached the top of the resin (1 mm thickness) add 1 ml of shaken SR5 solution to the capture column and allow it to gravity flow through it

31 Once the SR5 solution has reached the top of the resin shake solution SR6 and add 1 mL to the column

32 Immediately transfer the RNA capture column to a new clean 15 mL tube

33 Allow solution SR6 to flow through the column. Here wait until all the column is completely dry

34 Transfer the eluted RNA to a 2.2 mL Collection Tube (provided)

Shake solution SR4 and add 1 mL to the 2.2 mL collection tube. Close gently the lid and invert tube

- 35 to mix 10 times.
- 36 Incubate tube at -20 °C for 30 minutes
- 37 Centrifuge the 2.2 vials at 4 °C for 15 minutes at 13,000 x g

Note

After centrifugation the pellet may have a light brown color or white. If the final RNA yield is low may be barely visible and in this case is recommendable to place the vials in the centrifuge memorizing the position where the centrifugation force may deposit the pellet in order to have a proper final RNA resuspension.

- 38 Decant the supernatant
- 39 Quickly spin down the tubes to collect remaining drops on the tubes wall
- 40 Remove the supernatant using a smaller pipette (P20 or P100) without touching the pellet
- 41 Air dry the pellet for 5 minutes
- 42 Resuspend the RNA pellet in 40 µL of Solution SR7

- 43** Split the resuspended RNA in clean RNase free 200µL PCR tubes as follows.
Final Volume (FV) 40 µL

Tube 1 – 17 µL
Tube 2 – 17 µL
Tube 3 – ~ 6 µL

Note

Tube 3 will be used for RNA quantification and RIN assessment while tube number 1 will be used for the downstream DNase treatment. Tube 2 can be stored at -80°C as a back-up sample in case of problems during the downstream treatments.
If a low RNA yield is occurring (according to preliminary tests) resuspend the RNA pellet (step 42) in a 20 µL final volume and split the sample as follows:

*Tube 1 – 17 µL
Tube 2 – ~3 µL*

- 44** Place the PCR tubes on ice and proceed to step 45 or place the samples at -20 °C for short storage and -80 °C for longer

RNA yield assessment

- 45** Place the PCR tubes containing the eluted RNA on ice
- 46** Determine concentration of RNA using Qubit. Take 2-4 µL from tube #3

DNase treatment using DNase Max Kit

- 47** Thaw on ice the tube #1 containing 17µL total RNA extraction from step 43 [go to step #43](#)

48 Add 2µL of 10X DNase Buffer

49 Add 1 µL of DNase I enzyme (10 units)

50 Mix by gently pipetting

Safety information

Don't vortex the DNase I enzyme or the final mixture. It will denature the enzyme

51 Incubate the mixture, using a thermocycler, at 37 °C for 20 minutes and proceed to the next step, the DNase removal

52 Resuspend the DNase Removal resin by vortexing until the slurry is homogeneous

53 Add 5 µL of DNase removal resin to the mixture (step 51) go to step #51

54 Incubate for 10 minutes at Room temperature and check on the tube every 2 minutes with resuspending the resin by gently pipetting the solution

55 Centrifuge the tubes at 13.000 x g for 1 minute to pellet the resin

- 56 Transfer the supernatant containing the RNA target to a new clean 200µL tube without touching or removing the resin

Note

To avoid contaminant carryover pipette approximately 15µL

- 57 Determine the successful DNA removal procedure using Qubit or for higher detection limit perform a qPCR control. Use 1 to 5 µL from the previous step (56)

- 58 Determine concentration of RNA using Qubit. Use 1-2 µL

- 59 Proceed to the RIN assessment number or store the RNA at -80 °C

RNA quality assesment (RIN)

- 60 Allow all reagents to equilibrate to room temperature for 30 minutes before use and protect dye and dye mixture from light.

- 61 Set up the priming station for Bioanalyzer® chip adjusting the base plate in C position and sliding the syringe clip up to the top position

Safety information

Be sure of the clip position. Different position will lead to a negative output

- 62 Prepare the RNA ladder and the gel

Note

Ladder and gel preparation described in appendix 3 [⇒ go to step #135](#)

63 Vortex RNA dye concentrate (blue cap vial) for 10 seconds, spin down and add 1 μL of dye into a 65 μL aliquot of filtered gel

64 Vortex solution well. Spin tube at 13000g for 10 minutes at Room temperature

Note

Use prepared gel-dye mix within one day

65 Place a new RNA chip on the chip priming station

Note

Always insert the pipette tip to the bottom of the well when dispensing the liquid. Placing the pipette at the edge of the well may lead to poor results.

66

Pipette 9 μL of gel-dye mix in the well marked

67 Make sure that the plunger is positioned at 1 mL and then close the chip priming station

68 Press plunger until it is held by the clip

69 Wait for exactly 30 seconds then release clip

70 Wait for 5 seconds and slowly pull back plunger to 1 mL position

71 Open the chip priming station and pipette 9 μ L of gel-dye mix in the wells marked with

G

- 72 Pipette 9 μ L of the RNA conditioning solution (vial with white cap) into the well marked CS

- 73

Pipette 5 μ L of RNA marker (green cap vial) in all 11 sample wells and in the well marked

74

Pipette 1 μ L of the heat denatured and aliquoted ladder in the well marked

75

Pipette 1 μ L of sample in each of the 11 sample wells. Pipette 1 μ L of RNA Marker (green cap vial) in each unused sample well

Note

It is recommended to heat denature all RNA samples before use for 2 min and 70 °C (once) and keep them on ice

- 76** Place the chip in the Agilent 2100 Bioanalyzer and run the instrument within 5 min

Note

Set up the Bioanalyzer instrument prior loading the chip. Keeping loaded gel for more than 5 minutes will allow the liquid in it to evaporate leading to a poor or non-acceptable analysis. For a clear overview of the RNA integrity electropherogram output from the bioanalyzer see appendix 4 ➡ go to step #137

- 77** Place the chip horizontally in the IKA vortexer and vortex for 1 min at 2400rpm.

Safety information

DO NOT VORTEX FOR MORE THAN 1 MINUTE AND MORE THAN 2400rpm

Illumina library preparation

78

Note

This procedure was used for libraries without any enrichment or depletion of total RNA with RIN scores > 7 and a concentration of 100 ng purified RNA. RNA was prepared in 5 µL final volume of Nuclease-free water, to start with the first protocol step. The protocol is optimized for approximately 200 nt RNA inserts. To generate libraries with longer RNA insert sizes or lower RIN number, refer to Neb manufacturer protocol.

RNA Fragmentation and Priming

Note

Assemble all the reaction on ice using a nuclease-free tubes and in a dedicated clean PCR hood. The Illumina library preparation requires 7 steps using the thermocycler with different incubation times/cycles and temperatures; it is highly recommended to create a dedicated folder in the thermocycler with saved programs as suggested in the protocol in a chronological way. See appendix 5

⇒ go to step #138 to set up the thermocycler

Thaw the RNA samples from step 58 on ice ⇒ go to step #58

79 Perform dilutions, if necessary, to obtain a 100ng of total RNA in 5µL Final Volume using new clean 200µL PCR tubes

80 Set up the reaction (using the tubes from the previous step) as follows:

FRAGMENTATION AND PRIMING MIX	VOLUME
Purified mRNA (from step 78)	5 µL
• (lilac) NEBNext First Strand Synthesis Reaction Buffer	4 µL
• (lilac) Random Primers	1 µL
Total volume	10 µL

81 Set a 20 µL pipette to 5 µL and then pipette the entire volume up and down at least 10 times to

mix thoroughly. Perform a quick spin to collect all liquid from the sides of the tube

- 82 Place the sample in a thermocycler, upload and start program #1

Note

For program #1 see appendix 5 ➞ go to step #138

- 83 Once the incubation is ended, immediately transfer the tube to ice

- 84 Keeping the tubes on ice, assemble the following reaction:

FIRST STRAND SYNTHESIS REACTION	VOLUME
Fragmented and Primed RNA (Step 82)	10 µL
• (brown) NEBNext Strand Specificity Reagent	8 µL
• (lilac) NEBNext First Strand Synthesis Enzyme Mix	2 µL
Total volume	20 µL

- 85 Set a 20 µL pipette to 10 µL and then pipette the entire volume up and down at least 10 times to mix thoroughly. Perform a quick spin to collect all liquid from the sides of the tube

- 86 Place the sample in a thermocycler, upload and start program #2

Note

For program #2 see appendix 5 ➞ go to step #138

- 87 Once the incubation is ended, immediately transfer the tube to ice and proceed to next step

- 88 Keeping the tubes on ice, assemble the following reaction:

SECOND STRAND SYNTHESIS REACTION	VOLUME
First-Strand Synthesis Product (Step 86)	20 μ L
• (orange cap) NEBNext Second Strand Synthesis Reaction Buffer with dUTP Mix (10X)	8 μ L
• (orange cap) NEBNext Second Strand Synthesis Enzyme Mix	4 μ L
Nuclease-free Water	48 μ L
Total volume	80 μ L

- 89 Set a 100 μ L pipette to 40 μ L and then pipette the entire volume up and down at least 10 times to mix thoroughly. Perform a quick spin to collect all liquid from the sides of the tube

- 90 Place the sample in a thermocycler, upload and start program #3

Note

For program #3 see appendix 5 [➡ go to step #138](#)

- 91 Once the incubation is ended proceed to the first Purification step (91.1)

Purification of Double-stranded cDNA

91.1

Note

Before starting, prepare 80% ethanol; use a daily freshly prepared one. It is preferably to process no more than 6 samples at once; this will reduce the risk of over-dry the beads with a lower DNA target recovery. In this protocol the purification was performed using NEBNext Sample Purification Beads

Resuspend the NEBNext Sample Purification Beads by vortexing until the slurry is homogeneous

91.2 Add 144 μL (1.8X) of resuspended beads to the tubes from step 89 (second strand synthesis, ~80 μL) [⇒ go to step #89](#)

91.3 Mix well on a vortex mixer

91.4 Incubate for 5 minutes at Room temperature

91.5 Briefly spin the tube in a microcentrifuge to collect any sample on the sides of the tube

91.6 Place the tube on a magnetic rack for ~5 minutes to separate beads from the supernatant

91.7 Once the solution is clear, carefully remove and discard the supernatant without either removing or touching the beads

Safety information

Be careful with the beads, they retain the target DNA

91.8

Add 200 µL of 80% ethanol to the tube while in the magnetic rack

91.9

Incubate at Room temperature for 30 seconds and then carefully remove and discard the supernatant

91.10

Repeat from Step 91.8 once for a total of 2 washing steps

91.11

Briefly spin the tubes and put the tubes back on the magnetic rack

91.12

Remove the residual ethanol pipetting and air dry the beads for up to **5 minutes** while the tube is on the magnetic rack with lid open

Note

Do not over-dry the beads. Elute the samples when the beads are still dark brown and glossy looking, but when all visible liquid has evaporated. When the beads turn lighter brown and start to crack they are too dry

91.13

Remove the tube from the magnetic rack

91.14

Add 53µL of 0.1X TE Buffer (provided) to elute the DNA from the beads

91.15 Mix well on a vortex mixer

91.16 Quickly spin the tube and incubate for 2 minutes at Room temperature

91.17 Place the tube on the magnetic rack for ~5 minutes, until the solution is clear

91.18 Remove 50µL of the supernatant and transfer to a clean 200µL nuclease free PCR tube without disturbing the beads pellet

Note

If you need to stop, this is a safe stop point. Samples can be stored at -20°C

Illumina library preparation

92 Place the tube from step 91.18 on ice and assemble the following reaction:

END PREP REACTION	VOLUME
Second Strand Synthesis Product (Step 91.18)	50 µL
• (green cap) NEBNext Ultra II End Prep Reaction Buffer	7 µL
• (green cap) NEBNext Ultra II End Prep Enzyme Mix	3 µL
Total volume	60 µL

Note

Here is possible to prepare a mastermix. If a master mix is made, add 10µL of master mix to 50µL of cDNA for the End Prep reaction

- 93** Set a 100 µL pipette to 50 µL and then pipette the entire volume up and down at least 10 times to mix thoroughly. Perform a quick spin to collect all liquid from the sides of the tube

Note

It is important to mix well. The presence of a small amount of bubbles will not interfere

- 94** Place the sample in a thermocycler, upload and start program #4. Immediately proceed to the next step

Note

For program #4 see appendix 5 [go to step #138](#)

- 95**
- During the incubation, perform a 5-fold dilution of the (red) NEBNext Adaptor in Adaptor Dilution Buffer and keep the diluted adaptors on ice

Note

The NEBNext adaptor is provided in NEBNext oligos

- 96** Once both the Adaptor dilution and incubation are done, assemble the ligation reaction on ice by adding the following components

LIGATION REACTION	VOLUME
End Prepped DNA (Step 93)	60 µL
Diluted Adaptor (Step 95)	2.5 µL
• (red cap) NEBNext Ligation Enhancer	1 µL
• (red cap) NEBNext Ultra II Ligation Master Mix	30 µL
Total volume	93.5 µL

Note

Follow the order given. We do not recommend a premix

- 97** Set a 100 µL pipette to 80 µL and then pipette the entire volume up and down at least 10 times to mix thoroughly. Perform a quick spin to collect all liquid from the sides of the tube

Note

The NEBNext Ultra II Ligation Master Mix is very viscous. Care should be taken to ensure adequate mixing of the ligation reaction, as incomplete mixing will result in reduced ligation efficiency. The presence of a small amount of bubbles will not interfere with performance

- 98** Place the sample in a thermocycler, upload and start program #5

Note

For program #5 see appendix 5 [⇒ go to step #138](#)

99

Add 3 µL (blue cap vial) USER Enzyme to the ligation mixture after the incubation, resulting in total volume of 96.5 µL

100 Set a 100 μL pipette to 80 μL and then pipette the entire volume up and down at least 10 times to mix thoroughly. Perform a quick spin to collect all liquid from the sides of the tube

101 Place the sample in a thermocycler, upload and start program #6

Note

For program #6 see appendix 5 [⇒ go to step #138](#)

102 Once the incubation is ended proceed to the second Purification step

Purification of the Ligation Reaction

102.1 Resuspend the NEBNext Sample Purification Beads by vortexing until the slurry is homogeneous

102.2 Add 87 μL (0.9X) of resuspended beads to the tubes from step 100 (Ligation reaction, ~96.5 μL)

102.3 Mix well on a vortex mixer

102.4 Incubate for 10 minutes at Room temperature

102.5 Briefly spin the tube in a microcentrifuge to collect any sample on the sides of the tube

102.6 Place the tube on a magnetic rack for ~5 minutes to separate beads from the supernatant

102.7 Once the solution is clear, carefully remove and discard the supernatant without either removing or touching the beads

Safety information

Be careful with the beads, they retain the target DNA

102.8 Add 200 µL of 80% ethanol to the tube while in the magnetic rack

102.9 Incubate at room temperature for 30 seconds, and then carefully remove and discard the supernatant

102.10 Repeat from Step 102.8 once for a total of 2 washing steps

102.11 Briefly spin the tubes and put the tubes back on the magnetic rack

102.12 Remove the residual ethanol pipetting and air dry the beads for up to **5 minutes** while the tube is on the magnetic rack with lid open

Safety information

Do not over-dry the beads. Elute the samples when the beads are still dark brown and glossy looking, but when all visible liquid has evaporated. When the beads turn lighter brown and start to crack they are too dry

- 102.13** Remove the tube from the magnetic rack
- 102.14** Add 17µL of 0.1X TE Buffer (provided) to elute the DNA from the beads
- 102.15** Mix well on a vortex mixer
- 102.16** Quickly spin the tube and incubate for 2 minutes at Room temperature
- 102.17** Place the tube on the magnetic rack for ~5 minutes, until the solution is clear
- 102.18** Remove 15µL of the supernatant and transfer to a clean 200µL nuclease free PCR tube without disturbing the beads pellet

Note

If you need to stop, this is a safe stop point. Samples can be stored at -20°C

Illumina library preparation

103 Place the tube from step 102.18 on ice and assemble the following reaction:

Safety information

The following PCR Enrichment was performed specifically using NEBNext Multiplex Oligos for Illumina (Index Primers set 1) (NEB #E7335S) with the oligos concentration of 10 μ M. The PCR cycles are recommended using an initial RNA concentration of 100 ng; limit the PCR cycles to avoid overamplification with a secondary unwanted peak (~1000 bp) which will appear on the Bioanalyzer

PCR ENRICHMENT	VOLUME
Adaptor Ligated DNA (Step 102.18)	15 μ L
• (blue cap) NEBNext Ultra II Q5 Master Mix	25 μ L
Universal PCR Primer	5 μ L
Index (X) Primer	5 μ L
Total volume	50 μ L

Note

The Universal PCR Primer should be added to all the samples while the indexing should be unique for each sample. See appendix 6 [⇒ go to step #139](#) for an example of assembling reaction

104 Place the sample in a thermocycler, upload and start program #7

Note

For program #7 see appendix 5 [⇒ go to step #138](#)

105 Once the PCR amplification is ended proceed to the last Purification step

Purification of the PCR Reaction

105.1

Resuspend the NEBNext Sample Purification Beads by vortexing until the slurry is homogeneous

105.2

Add 45µL (0.9X) of resuspended beads to the tubes from step 100 (PCR reaction, ~50µL)

105.3

Mix well on a vortex mixer

105.4

Incubate for 5 minutes at Room temperature

105.5

Briefly spin the tube in a microcentrifuge to collect any sample on the sides of the tube

105.6

Place the tube on a magnetic rack for ~5 minutes to separate beads from the supernatant

105.7

Once the solution is clear, carefully remove and discard the supernatant without either removing or touching the beads

Safety information

Be careful with the beads, they retain the target DNA

- 105.8** Add 200µL of 80% ethanol to the tube while in the magnetic rack
- 105.9** Incubate at Room temperature for 30 seconds, and then carefully remove and discard the supernatant
- 105.10** Repeat from Step 105.8 once for a total of 2 washing steps
- 105.11** Briefly spin the tubes and put the tubes back on the magnetic rack
- 105.12** Remove the residual ethanol pipetting and air dry the beads for up to **5 minutes** while the tube is on the magnetic rack with lid open
- Safety information**

Do not over-dry the beads. Elute the samples when the beads are still dark brown and glossy looking, but when all visible liquid has evaporated. When the beads turn lighter brown and start to crack they are too dry
- 105.13** Remove the tube from the magnetic rack
- 105.14** Add 23µL of 0.1X TE Buffer (provided) to elute the DNA from the beads
- 105.15** Mix well on a vortex mixer

- 105.16** Quickly spin the tube and incubate for 2 minutes at Room temperature
- 105.17** Remove 20µL of the supernatant and transfer to a clean 200µL nuclease free PCR tube without disturbing the beads pellet
- 106** Determine concentration of DNA (Step 105.17) using Qubit. Use 1 µL
- 107** Store the DNA at -20°C or proceed to the library quality assessment on Bioanalyzer DNA chip

Library Average Fragment Length Assessment on an Agilen..

- 108** Allow all reagents to equilibrate to room temperature for 30 minutes before use and protect dye and dye mixture from light
- 109** Set up the priming station for Bioanalyzer® chip adjusting the base plate in C position and sliding the syringe clip up to the bottom position

Safety information

Be sure of the clip position. Different position will lead to a negative output

Preparing the Gel-Dye Mix

- 110** Add 15 µL of High Sensitivity DNA dye concentrate (blue) to a High Sensitivity DNA gel matrix vial (red)

111 Vortex solution well and spin down. Transfer to spin filter

112 Centrifuge at $2240\text{ g} \pm 20\%$ for 15 min. Store at $4\text{ }^{\circ}\text{C}$.

Safety information

Protect solution from light. Use prepared gel-dye mix within 6 weeks of preparation

Chip loading

Safety information

Always insert the pipette tip to the bottom of the well when dispensing the liquid. Placing the pipette at the edge of the well may lead to poor results

113 Place a new DNA chip on the chip priming station

114

Pipette $9\text{ }\mu\text{L}$ of gel-dye mix in the well marked

115 Make sure that the plunger is positioned at 1 mL and then close the chip priming station

116 Press plunger until it is held by the clip

117 Wait for exactly 60 seconds then release clip

118 Wait for 5 seconds and slowly pull back plunger to 1 mL position

119 Open the chip priming station and pipette 9 μL of gel-dye mix in the wells marked with

G

120

Pipette 5 µL of DNA marker (green cap vial) in all 11 sample wells and in the well marked

121

Pipette 1 µL of High Sensitivity DNA ladder (yellow cap vial) in the well marked

- 122** Pipette 1µL of sample in each of the 11 sample wells. Pipette 1µL of DNA Marker (green cap vial) in each unused sample well

- 123** Positionate the chip horizontally in the IKA vortexer and vortex for 1 min at 2400rpm

Safety information

DO NOT VORTEX FOR MORE THAN 1 MINUTE AND MORE THAN 2400rpm

- 124** Place the chip in the Agilent 2100 Bioanalyzer and run the instrument within 5 min

Note

Set up the Bioanalyzer instrument prior loading the chip. Keeping loaded gel for more than 5 minutes will allow the liquid in it to evaporate leading to a poor or non-acceptable analysis. For a clear overview of the DNA Average Fragment Length electropherogram output from the bioanalyzer see appendix 7 ➡ [go to step #140](#)

qPCR quantification of the library prepared for Illumina seq..

- 125** Place all the reagents of the KAPA Library thawing on ice

Note

If the kit is used for the first time, add the Primer Premix (10X) (1 mL) to the bottle of KAPA SYBR® FAST qPCR Master Mix (2X) (5 mL). Mix thoroughly using a vortex mixer

- 126** Prepare the appropriate library dilutions (using DNA dilution buffer). Depending on the expected concentration of the library, 1:1,000 – 1:100,000 dilutions may be appropriate

Note

All DNA Standards and library dilutions have to be assayed in triplicate. Each sample concentration should fall within the dynamic range of given standards. Check appendix 8 ➡ [go to step #141](#) *to see examples of dilution calculations*

- 127** Setup the following reaction on ice using 96-well 0.2 ml PCR Plates

qPCR MIX	VOLUME
KAPA SYBR FAST qPCR Master Mix (2X) + Primer Premix (10X)	12 µL
PCR-grade water	4 µL
Dilution of libraries	4 µL
Total volume	20 µL

128 Perform qPCR with the following cycling protocol:

Step	Temp.	Duration	Cycles
Initial denaturation	95°C	5 min	1
Denaturation	95°C	30 sec	35
Annealing/Extension / Data acquisition	60°C	45 sec	
Melt curve analysis	65 - 95°C		

129 Perform the results analysis to check the library quantification

Note

KAPABIOSYSTEMS provides a spreadsheet premade designed for the analysis of NGS library quantification data generated with the KAPA Library Quantification Kit for Illumina®. You can download it at the following link:

https://www.google.it/url?sa=t&rct=j&q=&esrc=s&source=web&cd=1&ved=2ahUKEwjAyPrV2s_hAhWxIYsKHW-mB9gQFjAAegQIARAC&url=https%3A%2F%2Fwww.kapabiosystems.com%2Fdocument%2Fkapa-library-quantification-data-analysis-template%2F%3Fdl%3D1&usg=AOvVaw0cKEI4Os4HF1F8a

130 Normalize all the DNA samples to the lowest amount of DNA

131 Pool the samples for the Illumina sequencing

132 Store the samples at -80°C

Appendix 1

133 G2 test

"G2 DNA/RNA Enhancer is convenient to use, when optimal DNA and or RNA extraction yield is required from especially clay. The primary function of G2 DNA/RNA Enhancer is to relieve inhibitory DNA-clay particle formations. G2 DNA/RNA Enhancer should be used in combination with either standardized extraction methods or commercial kits intended for DNA & RNA extraction from soil and clay" (Ampliqon).

Note

For this test were processed 500 mL and 1 mg of debris/soil for each sample with the addition of G2. According to protocol were added 2 mL of G2, to check the effect on the RNA extraction. A control without G2 was used.

Graphic representation of the higher RNA yield using G2

Appendix 2

- 134** Relative Centrifugal Force (**RCF**) is the term used to describe the amount of accelerative force applied to a sample in a centrifuge. **RCF** is measured in multiples of the standard acceleration due to gravity at the Earth's surface (x g). This is why **RCF** and "x g" are used interchangeably in centrifugation protocols.

$$\text{RCF or G} = 1.12 \times \text{Radius (mm)} \times (\text{rpm}/1000)^2$$

Appendix 3

135 Preparing the RNA Ladder

135.1 Spin the ladder down and pipette in an RNase-free vial

- 135.2** Heat denature the ladder for 2 min at **70 °C**
- 135.3** Immediately cool the vial on ice
- 135.4** Add 90µL of RNase-free water and mix thoroughly
- 135.5** Prepare aliquots in recommended 0.5 mL RNase-free vials with the required amount for typical daily use
- 135.6** Store aliquots at **-80 °C** . After initial heat denaturation, the frozen aliquots should not require repeated heat denaturation
- 135.7** Before use, thaw ladder aliquots on ice

Safety information

Avoid extensive warming

136 Preparing the Gel

- 136.1** Pipette 550 µL of RNA gel matrix (red) into a spin filter

136.2 Centrifuge at 1500g $\pm$ 20 % for 10 min at Room temperature

136.3 Aliquot 65 μ L filtered gel into 0.5 mL RNase-free microcentrifuge tubes. Use filtered gel within 4 weeks.
Store at 4 °C .

Appendix 4

137 In the following section can be understood how the bioanalyzer works and the regions evaluated to check the final quality score

Electropherogram detailing the regions that are indicative of RNA quality

Sample electropherograms used to assess the RNA Integrity Number (RIN) software. Samples range from intact (RIN 10), to degraded (RIN 2)

Appendix 5

138

PCR Tables

Program #1

Step	Temp.	Duration	Lid temperature
Incubation	94°C	15 min	105°C

Program #2

Step	Temp.	Duration	Cycles	Lid temperature
1	25°C	10 min	1	≥80°C
2	42°C	15 min		
3	70°C	15 min		
4	4°C	∞		

Program #3

Step	Temp.	Duration	Lid temperature
Incubation	94°C	60 min	≤40°C

Program #4

Step	Temp.	Duration	Cycles	Lid temperature
1	20°C	30 min	1	≥75°C
2	65°C	30 min		
4	4°C	∞		

Program #5

Step	Temp.	Duration	Lid temperature
Incubation	20°C	15 min	Off

Program #6

Step	Temp.	Duration	Lid temperature
Incubation	37°C	15 min	≥45°C

Program #7

Step	Temp.	Duration	Cycles
Initial denaturation	98°C	30 Sec	1
Denaturation	98°C	10 Sec	7
Annealing/Extension	65°C	75 sec	
Final Extension	65°C	5 min	1
Hold	4°C	∞	

Appendix 6

139

Examples of how to perform PCR enrichment (Indexing) with 4 samples:

Sample #1

<i>PCR Enrichment</i>	VOLUME
Adaptor Ligated DNA (Step 101.19)	15 μ L
• (blue cap) NEBNext Ultra II Q5 Master Mix	25 μ L
Universal PCR Primer	5 μ L
Index (1) Primer	5 μ L
Total volume	50 μ L

Sample #2

<i>PCR Enrichment</i>	VOLUME
Adaptor Ligated DNA (Step 101.19)	15 μ L
• (blue cap) NEBNext Ultra II Q5 Master Mix	25 μ L
Universal PCR Primer	5 μ L
Index (2) Primer	5 μ L
Total volume	50 μ L

Sample #3

<i>PCR Enrichment</i>	VOLUME
Adaptor Ligated DNA (Step 101.19)	15 μ L
• (blue cap) NEBNext Ultra II Q5 Master Mix	25 μ L
Universal PCR Primer	5 μ L
Index (3) Primer	5 μ L
Total volume	50 μ L

Sample #4

<i>PCR Enrichment</i>	VOLUME
Adaptor Ligated DNA (Step 101.19)	15 μ L
• (blue cap) NEBNext Ultra II Q5 Master Mix	25 μ L
Universal PCR Primer	5 μ L
Index (4) Primer	5 μ L
Total volume	50 μ L

Note

The only difference in the mixture is the Indexing choice (**Highlighted in Bold**). NebNext Multiplex Oligoes for Illumina (Index Primers Set 1) contains 12 index primers and they can all be used following the previous example for 12 samples.

Appendix 7

140 Assessment of DNA library fragment length average

Figure showing an example of correct RNA library size distribution on a Bioanalyzer. The electropherogram should show a narrow distribution with a peak size approximately 300 bp.

Note

If a peak at ~ 80 bp (primers) or 128 bp (adaptor-dimer) is visible in the Bioanalyzer traces; Bring up the sample volume (from Step 104.18) to 50 μ l with 0.1X TE buffer and repeat the SPRIselect Bead or NEBNext Sample Purification Bead Cleanup Step (Section 105.1). A dilution may be necessary for running on a Bioanalyzer High Sensitivity DNA Chip

Appendix 8

141 qPCR dilutions calculation

Prepare the appropriate library dilutions (using DNA dilution buffer). Depending on the expected concentration of the library, 1:1,000 – 1:100,000 dilutions may be appropriate.

It is recommended to perform a ten-fold serial dilution to obtain a more reliable dilution.

Follow the empiric calculation:

1 ng/μL from the library with a 300 bp length it's equal to say 3.5 nM in concentration

So, 1ng/μL = 3.5 nM in concentration

Practical calculation:

If we have a sample with 20 ng/μL from the library, multiply 20 ng/μL for 3.5 (conversion value) :

20 ng/μL x 3.5 = 70 nM

The value in μM has to be transformed in pM and diluted a proper number of times to fall in the following standard table provided by the KAPABiosynthesis.

DNA Standard 1	20 pM
DNA Standard 2	2 pM
DNA Standard 3	0.2 pM
DNA Standard 4	0.02 pM
DNA Standard 5	0.002 pM
DNA Standard 6	0.0002 pM

70 nM = 70.000 pM

Possibly we should dilute the samples to fall between standard 3 and 5.

70.000 pM has to be diluted at least 100.000 times in order to obtain a final concentration of 0.7 pM. In this case an additional 10-fold dilution of each sample is recommended.