SEP 21, 2020

🌐 Toxicity assay for mosquito larvicidal activity-Part2

avinash.kale¹

¹Domnic Colvin

UM-DAE Centre for Excellence in Basic Sciences

avinash.kale

OPEN ACCESS

DOI:

dx.doi.org/10.17504/protocols.io.bmjrk4m6

Document Citation: avinash.kale 2020. Toxicity assay for mosquito larvicidal activity-Part2. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.bmjrk4m6>

License: This is an open access document distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Created: Sep 21, 2020

Last Modified: Sep 21, 2020

DOCUMENT integer ID:

42321

The phenotypic characters of *C. quinquefasciatus* can be easily distinguished from the other species of mosquitoes. After collecting the dead larvae, they were identified phenotypically, with the help of an expert entomologist in the Zoonosis Department of the Haffkine Institute for Training, Research and Testing, using the keys of Barraud#. # Pocock, R. Fauna of British India. Mollusca. London 2, (1908).

The bioassay experiments were carried out at the Department of Zoonosis, Haffkine Institute for Training, Research and Testing, Mumbai, India. The Department is a Central Insecticide Board, Ghaziabad, New Delhi, India approved centre and is actively engaged in testing for Bio efficacy, persistency, repellence, testing against crawling & flying insects, mosquito repellence test on human skin, water resistance study and study of products for testing of Anti- Ant

and Anti Wasp activity as per WHO guidelines.

The mosquito larvae strains used here for various testing are the F1 generation of *Culex quinquefasciatus* in their third instar stage of development. These strains of mosquitoes have been used for reporting various mosquito repellents but their resistance or susceptibility to Bt is undetermined.