FEB 19, 2019

Tracking bleach synchronized worms

Priota Islam¹

¹Imperial College London

Behavioural Genomics

Priota Islam

Imperial College London

OPEN ACCESS

DOI:

dx.doi.org/10.17504/protocols.io.yb4fsqw

Protocol Citation: Priota Islam 2019. Tracking bleach synchronized worms.

protocols.io

<https://dx.doi.org/10.17504/protocols.io.yb4fsqw>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: Feb 19, 2019

Last Modified: Feb 19, 2019

PROTOCOL integer ID:

20572

Preparing Imaging plates

- 1
 - Prepare and pour low peptone NGM onto 35mm imaging plates
 - Put them in the cold room for at least 2 days before use
 - Seed imaging plates with 50ul of freshly made 1:10 solution (OP50:M9 solution) the day

before recording

- Leave on bench to dry with lid on overnight

Preparing worms

2

- Chunk the worms on 3 maintenance plates
- Bleach the worms after 2 days following the protocol for ***Bleach synchronization of *C. elegans****
- Refeed the starved L1s on 2 different NGM plates that are OP50 seeded, 72hr prior to the experiment day (Repeat for every strain to be tracked)

Day of recording

3

- Using a hair pick – pick 5 young adults onto the seeded imaging plates
- Put the plates on the tracker – agar side up with the lids off
- Wait 30min for the worms to acclimatise
- Record for 15min (See protocol ***Imaging on the multiworm tracker*** for detailed tracking instructions)