

Jan 29, 2019

Version 1

Transfection by Electroporation in Euplotes crassus V.1



In 1 collection

DOI

dx.doi.org/10.17504/protocols.io.xjgfkjw

Angela Piersanti¹

¹University of Camerino

Protist Research to Opti...



Angela Piersanti

University of Camerino

Create & collaborate more with a free account

Edit and publish protocols, collaborate in communities, share insights through comments, and track progress with run records.

Create free account





DOI: https://dx.doi.org/10.17504/protocols.io.xjgfkjw

Protocol Citation: Angela Piersanti 2019. Transfection by Electroporation in Euplotes crassus. **protocols.io** https://dx.doi.org/10.17504/protocols.io.xjgfkjw

License: This is an open access protocol distributed under the terms of the <u>Creative Commons Attribution License</u>, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited



Protocol status: Working

We use this protocol and it's working

Created: January 29, 2019

Last Modified: January 29, 2019

Protocol Integer ID: 19784

Keywords: Euplotes, Electroporation, euplotes crassus, transfection by electroporation, electroporation, transfection

Troubleshooting



- 1 2×104 Euplotes crassus cells were collected and resuspended in 0.3 M glucose solution (7% sea water and 93% of 0.3 M glucose solution).
- 2 Each round of transfection 250 µl of cells were used. 0.25 µg of Label IT® Plasmid Delivery Control Cy®3 (Mirus) were added alone or mixed with 2.5 µl of Lipofectamine® 2000 Transfection Reagent (Invitrogen) according to the supplier.
- 3 The sample was transferred to the 0.2 cm cuvette. Bio-Rad Gene Pulser was used. Conditions were set as follows: 0.2 kV, 25 μ FD, 100 Ω . Time constant around 1.2.
- 4 More than 50% of cells were viable after electroporation (few cells fused together), then cells were resuspended in 3 ml of sea water.
- 5 The plasmid was visible in the cytoplasm immediately after the electroporation by fluorescent microscope.