APR 23, 2019

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.z26f8he

External link:
<https://doi.org/10.1038/s41592-020-0796-x>

Protocol Citation: José A. Fernández Robledo 2019. Transfection of *Cryptocodinium cohnii* using labelled DNA. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.z26f8he>

MANUSCRIPT CITATION:
In progress, unpublished

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: In development
We are still developing and optimizing this protocol

Created: Apr 14, 2019

Last Modified: Apr 23, 2019

Transfection of *Cryptocodinium cohnii* using labelled DNA

Nature Methods

José A. Fernández Robledo¹

¹Bigelow Laboratory for Ocean Sciences

Protist Research to Optimize Tools in Genetics (PROT-G)

José A. Fernández Robledo

ABSTRACT

José A. Fernández Robledo
Nastasia J. Freyria

Bigelow Laboratory for Ocean Sciences

Keywords: Electroporation
and lipofection of
Cryptocodinium cohnii

Reagents

1 Liquid medium

- L1 + 10 gr/L glucose NPN (<https://ncma.bigelow.org/ccmp316#.XLOpDZNKi2w>)
- L1 medium with nitrogen limited (without yeast extract and glucose) to obtain cyst from *C. cohnii* swimming cells

Plates

- Prepare Petri dishes containing L1 medium bacteriological agar (1.5%)
- Prepare PEG 8000 solution (0.4%) with L1 medium, sterilized by using a filter (0.22µm)
- Pour 1 ml of L1-PEG8000 solution onto dry agar plates; alternatively, pour 12.5 ml of L1-PEG8000 solution onto dry agar plates
- Let the PEG800 solution infuse for 24 h
- Store at 4°C until use

Plasmid vectors

- Propagate *pPmMOE[MOE]:GFP-11* (Cold et al., 2016; Fernández Robledo et al., 2008) in JM109 for miniprep

pPmMOE[MOE]:GFP-11

- Linearize 20 µg with *NotI* (e.g., New England Biolabs® inc., Ipswich, MA, USA), clean the restriction digestion (e.g., GenElute® PCR Clean-up kit)
- Store at -20°C until use

DNA probe construction and DNA labeling

- *Use forward (5' -CCGCACATGTATGGTGAAGGCGAGGAGC- 3') the reverse (5' -CGTAGGACATGTCTTGTACAGCTCGTCCATGCCG- 3') primers targeting *pPmMOE[MOE]:GFP-11* (Fernández Robledo et al., 2008) to amplify a 739 bp DNA
- Clean the amplicon as above and label 1 µg using Ulysis™ Alexa Fluor® 488 Nucleic Acid

Labeling Kit (ThermoFisher Scientific) following the manufacturer's instructions

- Store labelled DNA were stored at -20°C until use

*Alternatively use any other set of primers/target DNA to generate the amplicon for labeling

References

Cold, E.R., Freyria, N.J., Martínez Martínez, J., Fernández Robledo, J.A., 2016. An agar-based method for plating marine protozoan parasites of the genus *Perkinsus*. PLoS One 11, e0155015.

Fernández Robledo, J.A., Lin, Z., Vasta, G.R., 2008. Transfection of the protozoan parasite *Perkinsus marinus*. Mol Biochem Parasitol 157, 44-53.

Cell Preparation

2 *Liquid culture*

- Incubate at 24°C in the dark with 125 rpm in a rotatory shaking incubator; the doubling time of *C. cohnii* cells under replete nitrogen conditions is 24-48 hours
- Harvest cells at log phase
- Centrifuge cells at 2,500 g for 10 min at room temperature
- Discard supernatant and maintain cells on ice until use

Spheroplasts induction

Cryptocodinium cohnii spheroplasts are derived from both swimming cells and cyst cells (Kwok et al., 2007; Pozdnyakov et al., 2014).

- Prepare PEG8000 (20%) (wt/vol) in L1 medium
- Resuspend *C. cohnii* swimming cells with 1 ml of 20% L1-PEG8000 solution and vortex for 10 min
- Spread 1 ml of *C. cohnii* swimming cells or cysts on Petri dishes containing L1 medium bacteriological agar and L1-PEG8000 solution (0.4%)
- Incubate for 2 days at 28°C
- Cover the colonies on the plate with L1 medium, pour off the medium
- Pour new L1 medium and retain the second elutant containing the spheroplasts
- Count spheroplasts using a haemocytometer (x3)

References

Kwok, A.C., Mak, C.C., Wong, F.T., Wong, J.T., 2007. Novel method for preparing spheroplasts from cells with an internal cellulosic cell wall. Eukaryot Cell 6, 563-567.

Pozdnyakov, I., Matantseva, O., Negulyaev, Y., Skarlato, S., 2014. Obtaining spheroplasts of armored dinoflagellates and first single-channel recordings of their ion channels using patch-clamping. Marine drugs 12, 4743-4755.

Electroporation

3

- Collect *Cryptocodinium cohnii* cells in log phase or spheroplasts (1.0 x10⁶ cells/mL)

- Electroporated with 5 µg (2.5 µg supercoiled, 2.5µg *NotI* linearized) of pPmMOE[MOE]GFP-11 using Amaxa®Cell Line Optimization Nucleofector™ solution V and program X-001 in a Nucleofector™ (Lonza)
- Recover cells from the cuvette and incubate in L1 medium
- As control for plasmid and electroporation, electroporate *P. marinus* PRA240 with pPmMOE[MOE]:GFP-11 as reported elsewhere (Fernández Robledo et al., 2008)
- Monitor cells for green fluorescence using standard FITC excitation/emission filters (488/507nm) under a transmitted-light fluorescence and/or confocal microscope

Lipofection

- 4
 - Lipofectamine® 3000 Reagent kit (ThermoFisher Scientific) in tissue culture plate 24-wells and 6-wells plates containing respectively 2- 3 mL of L1 medium
 - Use 15 µl of Lipofectamine and 1 µg of labeled DNA in 1.0×10^6 cells
 - Monitor cells for green fluorescence using standard FITC excitation/emission filters (488/507nm) under a transmitted-light fluorescence and/or confocal microscope

5

Expected result

Cryptocodinium cohnii viability after electroporation