VERSION 2

APR 30, 2018

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.ptsdnne

Protocol Citation: Carlos Helbig 2018. Transformation of *Vibrio natriegens*.
protocols.io
<https://dx.doi.org/10.17504/protocols.io.ptsdnne>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: In development
 We are still developing and optimizing this protocol

Created: Apr 27, 2018

Last Modified: Apr 30, 2018

PROTOCOL integer ID:
 11858

Transformation of *Vibrio natriegens* V.2

Carlos Helbig¹

¹iGEM

Carlos H

ABSTRACT

This protocol describes how to transform chemically competent *Vibrio natriegens* cells.

The protocol was described and published by Weinstock et al., 2016

BEFORE START INSTRUCTIONS

You need:

- Chemically competent *V. natriegens* cells
- BHI + V₂-Salts

10x V₂-Salts:

Chemical	Concentration	Molecular weight	Amount for 1L
NaCl	204mM	58.44g/mol	11.92176g
KCl	4.2mM	74.55g/mol	0.31311g
MgCl ₂	23.14mM	203.3g/mol	4.704362g

BHI:

For 1L medium 37g Brain heart infusion buillon

1 Defrost your competent cells on ice

- 2 Add your DNA template to the cells and mix gently
- 3 Incubate: 30min., on ice
- 4 Heatshock: 42°C, 45sec.
- 5 Incubate: 90sec., on ice
- 6 Add 1ml BHI + V₂-Salts (pre-heated 30°C) to the cells
- 7 Incubate: 30°C, 1min.
- 8 Recover: 30°C, 120min., 200rpm
- 9 Centrifuge: 3000g, 4min.

10 Plate the pellet on warm agar plates (30°C)

11 Incubate: