VERSION 2

SEP 27, 2021

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.bgtxjwpm

External link:
<https://www.neb.com/protocols/0001/01/01/transformation-protocol-for-bl21-de3-competent-cells-c2527>

Protocol Citation: New England Biolabs 2021. Transformation Protocol for BL21(DE3) Competent Cells (C2527I). **protocols.io** <https://dx.doi.org/10.17504/protocols.io.bgtxjwpm>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: May 25, 2020

Last Modified: Sep 27, 2021

Transformation Protocol for BL21(DE3) Competent Cells (C2527I) V.2

Forked from a private protocol

New England Biolabs¹

¹New England Biolabs

New England Biolabs (NEB)

Tech. support phone: +1(800)632-7799 email: info@neb.com

New England Biolabs

New England Biolabs

ABSTRACT

This transformation protocol is for the C2527I cells. (For the C2527H protocol, see [here](#).)

Keywords: T7 Expression Strain, comp cells, transforming for BL21(DE3)

Transformation Protocol Variables

Thawing: Cells are best thawed on ice and DNA added as soon as the last bit of ice in the tube disappears. Cells can also be thawed by hand, but warming above 0°C will decrease the transformation efficiency.

Incubation of DNA with Cells on Ice: For maximum transformation efficiency, cells and DNA should be incubated together on ice for 30 minutes. Expect a 2-fold loss in transformation efficiency for every 10 minutes you shorten this step.

Heat Shock: Both the temperature and the timing of the heat shock step are important and specific to the transformation volume and vessel. Using the transformation tube provided, 10 seconds at 42°C is optimal.

Outgrowth: Outgrowth at 37°C for 1 hour is best for cell recovery and for expression of antibiotic resistance. Expect a 2-fold loss in transformation efficiency for every 15 minutes you shorten this step. SOC gives 2-fold higher transformation efficiency than LB medium; and incubation with shaking or rotating the tube gives 2-fold higher transformation efficiency than incubation without shaking.

Plating: Selection plates can be used warm or cold, wet or dry without significantly affecting the transformation efficiency. However, warm, dry plates are easier to spread and allow for the most rapid colony formation.

Chemically competent *E. coli* cells suitable for transformation and protein expression.

Highlights

- Transformation efficiency: 1–5 x 10⁷cfu/μg pUC19 DNA
- T7 Expression Strain
- Deficient in proteases Lon and OmpT
- Resistant to phage T1 (*fhuA2*)
- B Strain
- Free of animal products

Genotype

fhuA2 [lon] ompT gal (λ DE3) [dcm] ΔhsdS

λ DE3 = λ sBamHI ΔEcoRI-B int:: (lacI::PlacUV5::T7 gene1) i21 Δnin5

MATERIALS

MATERIALS

BL21(DE3) Competent *E. coli* - 6x0.2 ml **New England Biolabs Catalog**
#C25271

SAFETY WARNINGS

Please refer to the Safety Data Sheets (SDS) for health and environmental hazards.

BEFORE START INSTRUCTIONS

Perform steps 2–9 in the tube provided.

- 1** Thaw a tube of BL21(DE3) Competent *E. coli* cells **On ice** until the last ice crystals disappear.

- 2** Mix gently and carefully pipette **50 μ L cells** into a transformation tube **On ice**.

- 3** Add **1 μ L** – **5 μ L** containing **1 pg** – **100 ng plasmid DNA** to the cell mixture.

- 4** Carefully flick the tube **4–5 times** to mix cells and DNA. **Do not vortex.**

- 5** Place the mixture **On ice** for **00:30:00**. Do not mix.

- 6** Heat shock at exactly **42 $^{\circ}$ C** for exactly **00:00:10**. Do not mix.

7 Place On ice for 00:05:00 . Do not mix.

8 Pipette 950 μL of Room temperature SOC into the mixture.

9 Place at 37 °C for 01:00:00 . Shake vigorously (250 rpm) or rotate.

10 Warm selection plates to 37 °C .

11 Mix the cells thoroughly by flicking the tube and inverting.

12 Perform several 10-fold serial dilutions in SOC.

13 Spread 50 μL – 100 μL of each dilution onto a selection plate and incubate Overnight at 37 °C .

Note

Alternatively, incubate at 30 °C for 24:00:00 – 36:00:00 or at 25 °C for 48:00:00 .