AUG 13, 2019

TRAP Assay

Eva Feldman¹

¹University of Michigan - Ann Arbor

Diabetic Complications Consortium

Tech. support email: rmcindoe@augusta.edu

Lili Liang

OPEN ACCESS

DOI:

dx.doi.org/10.17504/protocols.io.3sqgndw

External link:

<https://www.diacomp.org/shared/document.aspx?id=32&docType=Protocol>

Protocol Citation: Eva Feldman 2019. TRAP Assay.

protocols.io

<https://dx.doi.org/10.17504/protocols.io.3sqgndw>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: Jun 05, 2019

Last Modified: Aug 13, 2019

PROTOCOL integer ID:

24112

ABSTRACT

Summary:

The use of total radical-trapping antioxidant parameter (TRAP) has recently been proposed to explore the antioxidant property of a plasma sample. This assay is a measure of oxidative stress in the animals. This protocol describes the procedure used by the DiaComp to measure TRAP.

Diabetic Complications:

Cardiovascular

Retinopathy

Neuropathy

Nephropathy

Pediatric Endocrinology

Uropathy

Wound-Healing

Keywords: TRAP, cardiovascular, retinopathy, neuropathy, nephropathy, pediatric endocrinology, uropathy, wound-healing

MATERIALS

MATERIALS

⊗ 100 mM ABAP Sigma Aldrich Catalog #44,091-4

⊗ 30 mM PB pH 7.0 Contributed by users

⊗ Luminol Amersham Catalog #RPN2106

Reagent Preparation:

100mM ABAP*: Add 54.24 mg to 2 mL 30 mM PB.

Luminol: Mix 2 reagent $\frac{1}{2}$ & $\frac{1}{2}$..

BEFORE START INSTRUCTIONS

IMPORTANT: Be sure light shield is in place

Sample Preparation - DRG:

1. Remove 2 DRG from vial, cut in half and weigh. Do in duplicate.
2. Add 30 μ L 30mM PB and sonicate on 4 on ice.
3. Spin at maximum g's for 10 min at 4°C.
4. Remove sup and store on ice.

Performing the Assay:

2. 1. Using a **White Solid Bottom** plate, prepare plate by loading buffer for serial dilution. No PB in 1st well, 5 μ L PB in following 4 wells and 5 μ L in 3 control well. (Control is PB, Luminol and ABAP) (Dilutions, 1:1, 1:2, 1:4, 1:8, 1:16)
2. Prepare ABAP just prior to running assay by dissolving 54.2 mg ABAP in 2 mL PB. (Enough for 3 columns)
3. Load 5 μ L sample in wells 1 & 2. Mix the sample & PB in well 2 and remove 5 μ L and place in 3rd well with PB and so on. On last dilution discard 5 μ L.
4. Prepare Luminol in a 50 mL conical tube and add 200 μ L per well.

5. Place plate in Fluoroskan and add 60 μ L ABAP per well and press **START**.
6. Read every 30 seconds for 20 minutes.
7. When reading is done, Select Process>Organize. Choose the appropriate data to organize (usually Measure1), then click **OK**. This rearranges the data into columns.
8. Save organized data as an Excel file into the TRAP Assay data folder. Use the naming convention trXXXXXX.xls, where XXXXXX is the date in yymmdd format.

*ABAP = 2,2'-azobis(amidinopropane) dihydrochloride

*TRAP = **T**otal **R**adical-trapping **A**ntioxidant **P**arameter