MAY 10, 2018

TRIS buffer

Katy M
Monteith¹

¹University of Edinburgh

Vale Lab

Katy M Monteith
University of Edinburgh

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.p2vdqe6

Protocol Citation: Katy M Monteith 2018. TRIS buffer. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.p2vdqe6>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: May 10, 2018

Last Modified: May 10, 2018

PROTOCOL integer ID:
12085

Keywords: TRIS buffer, TRIS-HCl

ABSTRACT

This protocol describes how to make TRIS buffer (TRIS-HCl) at a 1M concentration and a pH 8.0

GUIDELINES

This protocol describes how to achieve a TRIS buffer at a pH 8.0. If a lower pH is required, i.e. pH 7.4, then add more hydrochloric acid until desired pH is reached

MATERIALS

MATERIALS

⊗ Tris Base, Molecular Biology Grade, 2,500g Promega Catalog #H5135

⊗ Hydrochloric acid Sigma Aldrich Catalog #H1758

SAFETY WARNINGS

! Read SDS for TRIS base and Hydrochloric acid prior to use.

1 Add 121.14g of TRIS base to 800 mL of triple distilled H₂O

- 2 Stir solution on magnetic plate using magnetic flea until TRIS base completely dissolved
- 3 Using a pH meter, adjust pH of solution to pH 8.0 by adding concentrated hydrochloric acid (HCl)
- 4 Add triple distilled H₂O to bring final volume of solution to 1 L
- 5 Autoclave