AUG 05, 2020

Trypsin digestion

avinash.kale¹

¹domnic colvin

UM-DAE Centre for Excellence in Basic Sciences

avinash.kale

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.bjdpki5n

Document Citation: avinash.kale 2020. Trypsin digestion.
protocols.io
<https://dx.doi.org/10.17504/protocols.io.bjdpki5n>

License: This is an open access document distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Created: Aug 05, 2020

Last Modified: Aug 05, 2020

DOCUMENT integer ID:
 40079

Trypsin digestion

In solution trypsin digestion of the TRIzol extracted protein sample was carried out by using Trypsin (Promega). Briefly, 15 μ L of Digestion Buffer (50 mM Ammonium bicarbonate) and 1.5 μ L of Reducing Buffer (100mM Dithiothreitol) were added to 10 μ L of the protein solution to the micro-centrifuge tube and the final volume was adjusted to 27 μ L with ultrapure water. The samples were incubated at 56 °C for 45 minutes and allowed to cool at room temperature. 3 μ L of freshly prepared alkylation buffer (100mM Iodoacetamide) was then added to the tube and kept for incubation in the dark at room temperature for 20 minutes. Finally, 10 μ L of Trypsin (20 μ g/ml) solution was added in the tube and kept for incubation at 37°C for 15 hours (overnight)(25).