



Jun 09, 2018

Two-minute walk test: reference equations for healthy adults and *the elderly in China*

DOI

dx.doi.org/10.17504/protocols.io.quqdwww

He Zou¹

¹Wenzhou Medical College



He Zou

Create & collaborate more with a free account

Edit and publish protocols, collaborate in communities, share insights through comments, and track progress with run records.

Create free account

OPEN  ACCESS



DOI: <https://dx.doi.org/10.17504/protocols.io.quqdwww>

Protocol Citation: He Zou 2018. Two-minute walk test: reference equations for healthy adults and the elderly in China. [protocols.io https://dx.doi.org/10.17504/protocols.io.quqdwww](https://dx.doi.org/10.17504/protocols.io.quqdwww)

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: June 09, 2018

Last Modified: June 09, 2018

Protocol Integer ID: 12912

Keywords: minute walk test, reference equations for healthy adult, elderly in china, healthy adult

Troubleshooting

Location

- 1 The two-minute walk test (2MWT) should be performed indoors, along a long, flat, straight, enclosed corridor with a hard surface that is seldom traveled. If the weather is comfortable, the test may be performed outdoors. The walking course must be 30 m in length.

Required equipment

- 2
 - (1). Countdown timer (or stopwatch)
 - (2). Mechanical lap counter
 - (3). Two small cones to mark the turnaround points
 - (4). A chair that can be easily moved along the walking course
 - (5). Worksheets on a clipboard
 - (6). A oximeter
 - (7). Sphygmomanometer
 - (8). Telephone

Subject preparation

- 3
 1. Comfortable clothing should be worn.
 2. Appropriate shoes for walking should be worn.
 3. A light meal is acceptable before early morning or early afternoon tests.
 4. Subjects should not have exercised vigorously within 2 hours of beginning the test.

Measurements

- 4
 1. Repeat testing should be performed about the same time of day to minimize intraday variability.
 2. A "warm-up" period before the test should not be performed.
 3. The subject should sit at rest in a chair, located near the starting position, for at least 10 minutes before the test starts. During this time, check for contraindications, measure pulse and blood pressure, and make sure that clothing and shoes are appropriate.
 4. Position the subject at the starting line. You should also stand near the starting line during the test. Each subject was asked to walk up and down the corridor as fast as possible in two minutes. As soon as the subject starts to walk, start the timer.
 5. The operator needed to prompted the subjects after 60 seconds using standardized encouragement ('you are doing well; you have one minutes left').
 6. Record the number of laps from the counter (or tick marks on the worksheet).
 7. Record the additional distance covered (the number of meters in the final partial lap) using the markers on the wall as distance guides. Calculate the total distance walked,



rounding to the nearest meter, and record it on the worksheet.

8. Congratulate the subject on good effort and offer a drink of water.