MAY 07, 2021

OPEN ACCESS

DOI:
[dx.doi.org/10.17504/protocols.io.bs2qnf5w](https://doi.org/10.17504/protocols.io.bs2qnf5w)

External link:
<https://doi.org/10.1371/journal.pone.0276248>

Protocol Citation: Lara J Carroll, Brijesh Tiwari, James F Curtin, Janith Wanigasekara 2021. U-251MG Spheroid generation using a scaffold based method protocol.
protocols.io
<https://dx.doi.org/10.17504/protocols.io.bs2qnf5w>

MANUSCRIPT CITATION:
 Wanigasekara J, Carroll LJ, Cullen PJ, Tiwari B, Curtin JF (2023) Three-Dimensional (3D) in vitro cell culture protocols to enhance glioblastoma research. PLOS ONE 18(2): e0276248.
<https://doi.org/10.1371/journal.pone.0276248>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

U-251MG Spheroid generation using a scaffold based method protocol

PLOS One Peer-reviewed method

Lara J Carroll¹, Brijesh Tiwari¹, James F Curtin¹,
 Janith
 Wanigasekara¹

¹Technological University Dublin

Technological University Dublin

PLOS ONE Lab Protocols

James F Curtin
 Technological University Dublin

ABSTRACT

3D cell culture is a technique that is used to grow cells in vitro that will mimic an in vivo environment. 3D cell models are a helpful learning tool for researchers to better understand disease mechanisms and to explore different therapeutic properties of drugs. 3D cell cultures can be developed using patient derived cancer cells. Once they have been grown, these 3D cells can be used to screen for small molecule drugs or for genetic modification in for analysis of disease pathways or to predict drug treatments toxicity or efficacy. 3D cell cultures are a big step towards the more ethical testing of drug toxicity and efficacy as they decrease the need to use animals in research as well as providing more reliable results as the cells used are of human physiology.

Cellusponge are 3D porous hydroxipropylcellulose scaffolds that are designed for use with cells that do not require specific ligands. As well as the standard non-coated cellusponge, there are two more of the same type of scaffold available for use that are made with two different coatings to allow for improved adaptation of different cell types, these are called Cellusponge-Gal and Cellusponge-Col. Cellusponge is a no-coating approach that is intended for use in the development of general soft tissue 3D culture. It has been used as soft matrix for 3D cell culture and 3D tumour model.

Protocol status: Working
We use this protocol and it's working

Created: Mar 04, 2021

Last Modified: May 07, 2021

PROTOCOL integer ID:
47888

Fig. 1: cellusponge 3D scaffolds. A) Placing a cellusponge disk in the middle of each well in a 24-well plate B) Three-dimensional porous hydroxipropylcellulose scaffold under 10x magnification C) cellusponge positioning after adding 500µL of complete media into each well in a 24-well plate

Fig. 2: U-251MG tumorspheroids formation in cellusponge 3D scaffolds. A) Tumorspheres formation after 1st day of incubation (Diameter - 22 μm), B) After 2nd day of incubation (Diameter - 49 μm), C) After 3rd day of incubation (Diameter - 70 μm), D) After 4th day of incubation (Diameter - 110 μm).

We used Bio-Byblos 9mm Cellusponge scaffold plates and adapted protocols published by:

1. ebersmedical.com. n.d. 3D Cellusponge Cell Seeding Protocol_Multi-well Plate Format Sponge. [online] Available at:
<https://ebersmedical.com/files/bbbmd/cellusponge_seeding_protocol.pdf> [Accessed 17 April 2021].
2. ebersmedical.com. n.d. Cellusponge Scaffolds. [online] Available at:
<<https://ebersmedical.com/tissue-engineering/scaffolds/cellusponge>> [Accessed

17 April 2021].

MATERIALS

- Complete medium (Dulbecco's Modified Eagle Medium + 10% V/V Fetal Bovine Serum + 1% Penicillin-Streptomycin (Sigma Aldrich)· Mammalian cells (U-251 MG human glioblastoma astrocytoma cells)
- Cellusponge 3D scaffolds (Sigma Alrich)
- Trypsin-EDTA solution(Sigma Aldrich)
- Phosphate Buffered Saline (PBS) 1X (pH41.7.4) [Sigma Aldrich]
- Trypan blue stain 0.4% (Thermo Fisher Scientific)
- Cell counter or cytometer
- Sterile reagent reservoirs
- Humidified incubator
- Multi-channel micropipette and tips
- 100-1000µl micropipette and tips
- Centrifuge with swinging bucket rotor
- Centrifuge tubes
- Bright-field microscope
- Cell culture flasks and dishes
- Serological pipettes

SAFETY WARNINGS

Laboratory coat and gloves must be worn at all times.

Spheroid Generation

- 1 Once U-251MG cell flasks are 70-80% confluent, medium from the flask is aspirated and U-251MG tumor cell monolayers re washed with 1X phosphate buffered saline (PBS; 5ml for a 25cm² or 4ml for a 75cm² flask), add 0.25 X Trypsin-EDTA cell dissociation enzyme (2ml for a 25cm² or 4ml for a 75cm² flask) and incubate cells at 37°C for 2-5 min.
- 2 Check cell detachment under a microscope and neutralize cell dissociation enzyme with complete growth medium. The 0.25 X Trypsin-EDTA reagent is neutralized using 4 volumes of complete medium.
- 3 Centrifuge U-251 MG single cell suspension at 5000rpm for 5 min, remove supernatant, tap the tube and re-suspend cell pellet in complete growth medium using a pipette. This should yield a single cell suspension without cell clusters.

- 4 Cells with a viability >90% can be used for spheroid generation. Mix 10µl of cells with 10µl of Trypan Blue and count cells using a haemocytometer.
- 5 For a 9mm scaffold, a cell suspension with the cell density of 5000k cells/ml must be obtained.
- 6 Slowly place a cellusponge disk in the middle of each well in a 24-well plate
- 7 Added 100µL of U-251MG cell suspension on top of the sponge for a 9mm disk and incubated it for 3 hours for U-251MG cells. This will remove any air bubbles within the cellusponge.
- 8 After incubation, 500µL of complete media was added to each well in a 24-well plate. Media should be added slowly along the edge of each well, not directly onto the cellusponge.
- 9 The 24-well plate cellusponge was then incubated overnight in an incubator (37°C, 5% CO₂, 95% humidity).
- 10 After overnight incubation, the seeded sponge is transferred into a new well plate, the media is replenished and the culture should be continued.
- 11 Tumor spheroid formation was observed within 3 days for U-251MG cell line.