MAY 10, 2019

U Mass - Chronic drug delivery

Jason Kim¹

¹University of Massachusetts

Mouse Metabolic Phenotyping Centers

Tech. support email: info@mmpc.org

Lili Liang

OPEN ACCESS

DOI:

dx.doi.org/10.17504/protocols.io.xuefnfte

External link:

<https://mmpc.org/shared/document.aspx?id=151&docType=Protocol>

Protocol Citation: Jason Kim 2019. U Mass - Chronic drug delivery. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.xuefnfte>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Feb 06, 2019

Last Modified: May 10, 2019

PROTOCOL integer ID:
20070

ABSTRACT

Summary:

A subcutaneous or intraperitoneal implantation of Alzet osmotic pump is used to chronically administer selected drug in mice. Chronic drug delivery may be used to examine intermediate to long-term effects of selected drug on obesity, insulin resistance, and metabolism.

MATERIALS

MATERIALS

Osmotic pump **Alzet Catalog #1007D**

Reagent Preparation:

Reagent 1: Drug or placebo containing osmotic pump

Reagents and Materials:

1. Alzet osmotic pump

Procedure:

1. Prepare drug or placebo solution based on dosage.
2. Load solution into a syringe.
3. Hold the pump with sterilized tweezers and slowly fill the solution (drug/placebo).
4. Close the hole.
5. For immediate delivery of compound upon subcutaneous implantation, osmotic pumps may be warmed by submerging them in warm water immediately prior to surgery.

Keywords: Chronic drug delivery, obesity, insulin resistance, metabolism

- 1 Anesthetize mice with an intraperitoneal injection of ketamine (100 mg/kg body weight) and xylazine (10 mg/kg body weight).
- 2 Shave hair at the incision site on the back.
- 3 Make an incision (~0.5 cm) using sterilized scalpel between the scapulae.
- 4 Subcutaneously insert an Alzet mouse osmotic pump containing drug or placebo.
- 5 Suture or close the incision site using sterilized staples.
- 6 Administer ketoprofen to minimize pain and house mice individually.