MAY 10, 2019

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.xxafpie

External link:
<https://mmpc.org/shared/document.aspx?id=141&docType=Protocol>

Protocol Citation: Jason Kim 2019. U Mass - Glucose Tolerance Test. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.xxafpie>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: Feb 08, 2019

Last Modified: May 10, 2019

PROTOCOL integer ID:
 20162

U Mass - Glucose Tolerance Test

Jason Kim¹

¹University of Massachusetts

Mouse Metabolic Phenotyping Centers
 Tech. support email: info@mmpc.org

Lili Liang

ABSTRACT

Summary:

Glucose tolerance test measures systemic clearance of glucose following an intraperitoneal bolus injection of 20% dextrose. This experiment measures insulin sensitivity in awake mice assuming that there are no alterations in the animal's pancreatic β -cell function and insulin secretion. Insulin sensitivity is altered in obese mice.

MATERIALS

MATERIALS

⊗ 20% Dextrose injection USP Hospira(Pfizer) Catalog #NDC0409-7935-19

Note:

[Hospira](#), [RRID:SCR_003985](#)

Keywords: Glucose
Tolerance Test, obesity

- 1 Mice may be fasted overnight (~15 hours) or for 5 hours prior to the start of experiment.
- 2 Collect plasma sample (10 μ l) before the start of experiment (basal-0 min) to measure basal glucose levels.
- 3 Administer intraperitoneal injection of 20% dextrose (1 or 2 g/kg body weight) using an insulin syringe.
- 4 Collect plasma samples (10 μ l) at 10, 20, 30, 60, 90, and 120 min following injection to measure circulating glucose concentrations.
- 5 For data analysis, plasma glucose levels vs. time after injection are plotted, and area under curve may be calculated to estimate insulin sensitivity.
- 6 Area under curve of glucose tolerance test may be inversely correlated with insulin sensitivity assuming unaffected insulin secretion and pancreatic β -cell function in mice.