May 10, 2019

O U Mass - Glucose Tolerance Test

DOI

dx.doi.org/10.17504/protocols.io.xxafpie

Jason Kim¹

¹University of Massachusetts

Mouse Metabolic Phenotyping Centers Tech. support email: info@mmpc.org

෯ Lili Liang





DOI: dx.doi.org/10.17504/protocols.io.xxafpie

External link: <u>https://mmpc.org/shared/document.aspx?id=141&docType=Protocol</u>

Protocol Citation: Jason Kim 2019. U Mass - Glucose Tolerance Test. protocols.io <u>https://dx.doi.org/10.17504/protocols.io.xxafpie</u>

License: This is an open access protocol distributed under the terms of the **<u>Creative Commons Attribution License</u>**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working We use this protocol and it's working

Created: February 08, 2019

Last Modified: May 10, 2019

Protocol Integer ID: 20162

Keywords: Glucose Tolerance Test, obesity



Abstract

Summary:

Glucose tolerance test measures systemic clearance of glucose following an intraperitoneal bolus injection of 20% dextrose. This experiment measures insulin sensitivity in awake mice assuming that there are no alterations in the animal's pancreatic β -cell function and insulin secretion. Insulin sensitivity is altered in obese mice.

Materials

MATERIALS

20% Dextrose injection USP **Pfizer (Hospira) Catalog #**NDC0409-7935-19

Note:

Hospira, RRID:SCR_003985

- 1 Mice may be fasted overnight (~15 hours) or for 5 hours prior to the start of experiment.
- 2 Collect plasma sample (10 μl) before the start of experiment (basal-0 min) to measure basal glucose levels.
- 3 Administer intraperitoneal injection of 20% dextrose (1 or 2 g/kg body weight) using an insulin syringe.
- 4 Collect plasma samples (10 μl) at 10, 20, 30, 60, 90, and 120 min following injection to measure circulating glucose concentrations.
- 5 For data analysis, plasma glucose levels vs. time after injection are plotted, and areaunder-curve may be calculated to estimate insulin sensitivity.
- 6 Area-under-curve of glucose tolerance test may be inversely correlated with insulin sensitivity assuming unaffected insulin secretion and pancreatic β-cell function in mice.