MAY 10, 2019

U Mass - Hepatic gluconeogenesis

Jason Kim¹

¹University of Massachusetts

Mouse Metabolic Phenotyping Centers

Tech. support email: info@mmpc.org

Lili Liang

OPEN ACCESS

DOI:

dx.doi.org/10.17504/protocols.io.xy7fpzn

External link:

<https://mmpc.org/shared/document.aspx?id=144&docType=Protocol>

Protocol Citation: Jason Kim 2019. U Mass - Hepatic gluconeogenesis.

protocols.io

<https://dx.doi.org/10.17504/protocols.io.xy7fpzn>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Feb 11, 2019

Last Modified: May 10, 2019

PROTOCOL integer ID:
20223

ABSTRACT

Summary:

Hepatic gluconeogenesis is estimated using pyruvate tolerance test that measures systemic elevation of glucose partly derived from pyruvate and hepatic gluconeogenesis following an intraperitoneal bolus injection of pyruvate in awake mice. Hepatic gluconeogenesis is affected by obesity and regulates glucose homeostasis.

MATERIALS

MATERIALS

Sodium Pyruvate **Sigma Aldrich Catalog #P5280**

Reagent Preparation:

Reagent 1: 10 % Pyruvate in PBS

Reagents and Materials: Sodium pyruvate, PBS (phosphate buffered saline)

Procedure: Dissolve 1 g of sodium pyruvate in 10 ml PBS

Note:

Sigma-Aldrich, [RRID:SCR_008988](#)

Keywords: Hepatic
gluconeogenesis, obesity

- 1 Mice are fasted overnight (~15 hours) prior to the start of experiment.
- 2 Collect plasma sample (10 μ l) before the start of experiment (basal-0 min) to measure basal glucose levels.
- 3 Administer intraperitoneal injection of pyruvate (1 g/kg body weight) using an insulin syringe.
- 4 Collect plasma samples (10 μ l) at 10, 20, 30, 45, 60, 90, and 120 min following pyruvate injection to measure circulating glucose concentrations.
- 5 For data analysis, plasma glucose levels vs. time after pyruvate injection are plotted, and area-under-curve may be calculated to estimate hepatic gluconeogenesis.