MAY 10, 2019

U Mass - Insulin tolerance test

Jason Kim¹

¹University of Massachusetts

Mouse Metabolic Phenotyping Centers

Tech. support email: info@mmpc.org

Lili Liang

OPEN ACCESS

DOI:

dx.doi.org/10.17504/protocols.io.x2xfqfn

External link:

<http://mmpc.org/shared/document.aspx?id=143&docType=Protocol>

Protocol Citation: Jason Kim 2019. U Mass - Insulin tolerance test. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.x2xfqfn>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Feb 12, 2019

Last Modified: May 10, 2019

PROTOCOL integer ID:
20279

ABSTRACT

Summary:

Insulin tolerance test measures systemic clearance of glucose following an intraperitoneal bolus injection of a physiological dose of insulin. This experiment measures insulin sensitivity in awake mice assuming that there are no alterations in the animal's counterregulatory response. Insulin sensitivity is altered in obese mice.

MATERIALS

MATERIALS

Insulin **Novolin Catalog #Regular human insulin, U-100**

Keywords: Insulin tolerance test, obese mice

- 1 Mice are fasted for 5 hours prior to the start of experiment.
- 2 Collect plasma sample (20 μ l) before the start of experiment (basal-0 min) to measure basal insulin and glucose levels.
- 3 Administer intraperitoneal injection of insulin (0.5 or 0.75 unit/kg body weight) using an insulin syringe.
- 4 Collect plasma samples (10 μ l) at 10, 20, 30, 60, 90 and 120 min for the measurement of glucose concentrations.
- 5 For data analysis, plasma glucose levels vs. time after insulin injection are plotted, and area-under-curve may be calculated to estimate insulin sensitivity.