

May 10, 2019

U Mass - Organ-specific glucose uptake

DOI

dx.doi.org/10.17504/protocols.io.x4cfqsw



Jason Kim¹

¹University of Massachusetts

Mouse Metabolic Phenotyping Centers
Tech. support email: info@mmpc.org



Lili Liang

OPEN  ACCESS



DOI: dx.doi.org/10.17504/protocols.io.x4cfqsw

External link: <https://mmpc.org/shared/document.aspx?id=138&docType=Protocol>

Protocol Citation: Jason Kim 2019. U Mass - Organ-specific glucose uptake. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.x4cfqsw>

License: This is an open access protocol distributed under the terms of the **Creative Commons Attribution License**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: February 13, 2019

Last Modified: May 10, 2019

Protocol Integer ID: 20324

Keywords: Glucose uptake, obese mice, glucose uptake in individual organ, specific glucose uptake summary, glucose uptake, glucose metabolism, glucose, reduced glucose metabolism, insulin resistance, metabolite levels in select tissue, labeled metabolite level, individual organ, select tissue, organ

Abstract

Summary:

Glucose uptake in individual organs can be measured using a bolus injection of 2-deoxy-D-[1-¹⁴C] glucose, a non-metabolizable glucose analog, and by determining labeled metabolite levels in select tissues. Insulin resistance is characterized by reduced glucose metabolism and develops in obese mice.

Materials

MATERIALS

⊗ Poly-prep columns prefilled with AG 1-X8 resin **Bio-Rad Laboratories Catalog #731-6211**

⊗ 0.2 M formic acid **Merck MilliporeSigma (Sigma-Aldrich) Catalog #F0507**

⊗ 0.5 M ammonium acetate **Merck MilliporeSigma (Sigma-Aldrich) Catalog #A1542**

Reagent Preparation:

Reagent 1: 0.2 M formic acid/0.5 M ammonium acetate

Reagents and Materials: formic acid, ammonium acetate, deionized water

Procedure:

1. Prepare 900 ml of dH₂O, and add 7.69 ml of formic acid.
2. Add 38.84 g of ammonium acetate, and adjust pH to 4.9±0.05 using dH₂O.
3. Add dH₂O to make a final solution volume of 1,000 ml.

Note:

Bio-Rad Laboratories [RRID:SCR_008426](#)

Sigma-Aldrich [RRID:SCR_008988](#)

- 1 Survival surgery is performed to establish a chronic indwelling catheter at 5~6 days prior to experiment for intravenous infusion. (refer to M1023: Surgery-jugular vein cannulation)
- 2 Mice are fasted overnight (~15 hours) or for 5 hours prior to the start of experiment.
- 3 Place a mouse in a rat-size restrainer with its tail tape-tethered at one end.
- 4 Administer an intravenous bolus injection of 10 μ Ci of 2-deoxy-D-[1- 14 C] glucose (2-[14 C]DG) in awake mice. Alternatively, intraperitoneal injection of 10 μ Ci of 2-[14 C]DG may be used in awake mice.
- 5 After 30 min, rapidly freeze-clamp the tissues in liquid N₂, and store tissue samples in -80°C freezer for biochemical assay.
- 6 Biochemical assay is conducted using frozen tissue samples (e.g., skeletal muscle, adipose tissue, heart) to measure tissue levels of 2-[14 C]DG-6-phosphate.
 - a) Prepare a heat block set to ~ 100°C.
 - b) Prepare anion-exchange columns by washing with 5 ml of dH₂O.
 - c) Homogenize 50–100 mg of frozen tissue samples by adding ten times the volume of dH₂O (50 mg of tissue in 500 μ l of dH₂O) in glass tubes using a tissue homogenizer.
 - d) Following homogenization, place the glass tubes in the heat block for 10 min, vortex for 2 sec, and then cool to room temperature.
 - e) Transfer the homogenized samples to microcentrifuge tubes using transfer pipettes and centrifuge at 16,000 \times g for 5 min.
 - f) Add 33 μ l of homogenate (supernatant) to 467 μ l dH₂O in a scintillation vial labeled "total" sample.
 - g) Add 5 ml of scintillation cocktail, vortex, and count the samples for 14 C using a liquid scintillation counter (total 14 C samples).
 - h) Transfer 333 μ l of homogenate (supernatant) to the anionexchange columns for the separation of 2-[14 C]DG-6-P from 2-[14 C] DG.

- i) Wash the columns with 2 ml of dH₂O three times and collect the samples into a scintillation vial labeled "wash" sample.
- j) Vortex the "wash" samples, and transfer 500 µl of "wash" samples to another set of scintillation vials to be counted for ¹⁴C using a liquid scintillation counter (wash samples containing 2-[¹⁴C] DG).
- k) Elute the columns with 2 ml of 0.2 M formic acid/0.5 M ammonium acetate three times, and collect the samples into a scintillation vial labeled "eluate" sample.
- l) Vortex the "eluate" samples, and transfer 500 µl of "eluate" samples to another set of scintillation vials to be counted for ¹⁴C using a liquid scintillation counter (eluate samples containing 2-[¹⁴C] DG-6-P).
- 7 The rate of glucose uptake in individual organs is determined using 2-[¹⁴C] DG. 2-[¹⁴C] DG is taken up by cells, phosphorylated by glucokinase to become 2-[¹⁴C] DG-6-P, and not further metabolized. Thus, organ-specific accumulation and level of 2-[¹⁴C] DG-6-P following a bolus injection of 2-[¹⁴C] DG reflect glucose uptake in individual organs.