VERSION 2

AUG 12, 2019

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.56ig9ce

External link:
<https://mmpc.org/shared/document.aspx?id=326&docType=Protocol>

Protocol Citation: Mark Kelly, Timothy P. Fitzgibbons 2019. U Mass - Transverse Aortic Constriction.
protocols.io
<https://dx.doi.org/10.17504/protocols.io.56ig9ce>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: Aug 05, 2019

Last Modified: Aug 12, 2019

U Mass - Transverse Aortic Constriction V.2

Mark Kelly¹, Timothy P. Fitzgibbons¹

¹University of Massachusetts Medical School

Mouse Metabolic Phenotyping Centers
 Tech. support email: info@mmpc.org

Lili Liang

ABSTRACT

Summary:

This is a Mouse model of Transverse aortic constriction (TAC). A commonly used experimental model for pressure overload-induced cardiac hypertrophy and heart failure. Initially leading to compensated hypertrophy of the heart, which often is associated with a temporary enhancement of cardiac contractility. Resulting in cardiac dilatation and heart failure.

MATERIALS

MATERIALS

- ☒ Ketamine **Patterson Veterinary Catalog # 07-803-6637**
- ☒ Xylazine **Patterson Veterinary Catalog #07-869-6707**
- ☒ Isoflurane 1-3% **Patterson Veterinary Catalog #07-893-1389**
- ☒ Buprenorphine **Patterson Veterinary Catalog #07-891-9756**
- ☒ Meloxicam **Patterson Veterinary Catalog #07-893-1368**
- ☒ 6-0 Prolene suture **Patterson Veterinary Catalog #07-824-3204**
- ☒ 5-0 Silk **Patterson Veterinary Catalog #07-824-1980**

Keywords: Transverse Aortic
Constriction (TAC)

1 Expected procedure duration:

20-40 minutes

2 Adequacy or depth of anesthesia is monitored by:

Respiratory Rate and Toe Pinch

3 Frequency of anesthesia depth assessment:

At the start of surgical procedure, a toe or ear pinch can be used to assess the depth of anesthesia. Visual monitoring should be performed throughout the procedures, as well as toe/ear pinches.

Deviations from expected behavior Should be noted.

4 Anesthesia Regimen:

Ketamine (80-100mg/kg), Xylazine (5-20mg/kg) or Isoflurane 1-3%

5 Pre-surgical Analgesics:

Approximately 30 minutes prior to undergoing the surgical procedure, mice receive an S.C. injection of Buprenorphine (0.05mg/kg) and Meloxicam (5mg/kg).

6 Surgical prep:

Aseptic technique will be maintained by:
Clipping/shaving fur around incision site, Sterile Instruments.

Isoflurane, heating pad, forceps, scissors, needle driver, eye ointment, 5-0 Prolene, 7-0 Prolene suture, 20g iv catheter sheath, 1 ml-syringes, 0.9% NaCl, Sterile gloves, Povidone-iodine, 70% ETOH

7 Transverse aortic constriction (TAC) Procedure:

- (1). Anesthetize the mouse and ensure depth of anesthesia with a toe pinch.
- (2). The mouse is positioned supine on a heating pad.
- (3). Shave the ventral neck and left parasternal area.
- (4). Prep the surgical field with 70% isopropanol as well as Betadine solution and drape.
- (5). Perform a longitudinal midline cervical incision from supra-sternal notch to the mid-chest to expose the sternum.
- (6). Retract the thyroid by passing through a 6-0 Prolene stay suture and tape it to the sterile working area.
- (7). Bluntly dissect the pre-tracheal muscles with micro-surgical forceps to expose the trachea, continue blunt dissection under the pre-tracheal muscles and behind the sternum to move the pleura away, gently elevate the right supra-clavicular muscles with forceps and carefully perform an 3-4 mm upper partial sternotomy by cutting the upper sternum, Using a retractor or place a 6-0 Prolene stay suture through each side of the sternotomy and tape it to the sterile working area.
- (8). Move aside the pre-tracheal muscles, mediastinal fat and thymus to visualize the aortic arch use caution not damage the parietal pleura to prevent pneumothorax development.
- (9). Blunt dissect the soft tissue under the aortic arch, prepare a tunnel in the soft tissue under the aortic arch, pass a segment of 5-0 silk suture under the aorta between the innominate and left common carotid arteries.
- (10). Place a 27g cannula needle alongside the aorta and tie the suture snugly around the needle and the aorta between the right innominate and left common carotid arteries. When complete, quickly but gently remove the needle to achieve ~0.4 mm diameter narrowing and a 65-70% aortic constriction.
- (11). Remove all stay suture and close the surgical site using 6-0 Prolene sutures.
- (12). One day following the TAC procedure, the mouse is subjected to Doppler echocardiography to determine the degree of stenosis induced by the ligation.

8 Post-procedure Analgesics:

Buprenorphine (0.05mg/kg) every 12 hours, for 72 hours post-op.
Meloxicam (5mg/kg) every 24 hours, for 72 hours post-op

9 Post-procedure Monitoring:

Mice are monitored 2x daily for the first 72 hours after the surgery. Thereafter, mice are monitored at least 3x per week.