MAY 13, 2019

U Michigan - Retinal Microstructure Imaging OCT

David A. Antonetti¹

¹University of Michigan - Ann Arbor

Mouse Metabolic Phenotyping Centers
Tech. support email: info@mmpc.org

Lili Liang

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.yabfsan

External link:
<https://mmpc.org/shared/document.aspx?id=308&docType=Protocol>

Protocol Citation: David A. Antonetti 2019. U Michigan - Retinal Microstructure Imaging OCT. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.yabfsan>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Feb 18, 2019

Last Modified: May 13, 2019

PROTOCOL integer ID:
20515

ABSTRACT

Summary:

A spectral domain ophthalmic imaging system based on the technology of optical coherence tomography (OCT) will be used to acquire the microstructure image of retina in live animals. Like the ultrasound, but uses light waves instead of sound waves, the light backscattered from within a sample is processed to develop a high-resolution, depth-resolved image suitable for analyzing internal microstructure in vivo.

MATERIALS

MATERIALS

⊗ Ketamine **Hospira(Pfizer) Catalog #0409-2051-05**

⊗ Xylazine **VetOne Catalog #510004**

⊗ Tropicamide Ophthalmic Solution 1% **Bausch & Lomb**

⊗ Phenylephrine Hydrochloride Ophthalmic Solution 2.5% **Akorn**

⊗ Propylene Glycol Eye Drop (Systane) **Alcon**

⊗ 0.3 c.c. insulin syringe (31-gauge x 5/16") **BD Biosciences Catalog #328440**

Note:

Hospira, [RRID:SCR_003985](#)

BD Biosciences, [RRID:SCR_013311](#)

Keywords: optical coherence tomography (OCT), internal microstructure

- 1 The eyes of conscious animals are dilated by first applying a small drops (~10 µl) of tropicamide (1%) followed by a small drop of phenylephrine (2.5%) 2 to 3 minutes later
- 2 Animal is then sedated with ketamine and xylazine
- 3 After sedation, the corneas are kept moist with the wetting agents (Systane)
- 4 Animal is then placed on a platform enabling three-dimensional fine movements
- 5 The camera lens is adjusted to aim at the eye to acquire images without physical contact
- 6 Volume or radial scan (1.4 x 1.4 mm) is obtained from each eye with a Bioptigen Envisu R2200 preclinical spectral domain optical coherence tomography (SD-OCT) system
- 7 After imaging procedure, the animal is allowed to recover from the anesthesia
- 8 Bioptigen Diver software is used to analyze retinal structure from the image obtained