



Aug 17, 2023

Version 2

U54 SCENT Pediatric Colonoscopy Tissue Collection Procedure V.2



DOI

dx.doi.org/10.17504/protocols.io.yxmvm3oznl3p/v2

Mary Jordan*¹

¹Duke BioRepository & Precision Pathology Center, Duke University Hospital, Durham, NC, USA

Cellular Senescence Net...



Valerie Bekker

Duke

Create & collaborate more with a free account

Edit and publish protocols, collaborate in communities, share insights through comments, and track progress with run records.

Create free account

OPEN  ACCESS



DOI: <https://dx.doi.org/10.17504/protocols.io.yxmvm3oznl3p/v2>

Protocol Citation: Mary Jordan* 2023. U54 SCENT Pediatric Colonoscopy Tissue Collection Procedure. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.yxmvm3oznl3p/v2> Version created by **Valerie Bekker**



License: This is an open access protocol distributed under the terms of the **Creative Commons Attribution License**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: August 17, 2023

Last Modified: August 17, 2023

Protocol Integer ID: 86610

Keywords: u54 scent pediatric colonoscopy tissue collection procedure, acquisition for pediatric colonoscopy colon specimen, pediatric colonoscopy colon specimen, u54 scent, department of pathology, precision pathology center, colon, duke university hospital through the biorepository, specimen, pathology, brpc, duke university hospital, biorepository

Abstract

This document outlines the required criteria and acquisition for pediatric colonoscopy colon specimens collected at Duke University Hospital through the Biorepository and Precision Pathology Center (BRPC) in the Department of Pathology.

Troubleshooting

Inclusion and Exclusion Criteria

1 Inclusion Criteria:

- Healthy patients age 5-18 years of age presenting for outpatient colonoscopy.
- Screening may wish to focus on the following indications: abdominal pain, constipation, diarrhea, bloating/gas, patients with normal labs and/or imaging.

Exclusion Criteria:

- Systemic Illness such as diabetes is a relative exclusion but may be hard to avoid
 - Autoimmune disease such as SLE (lupus) or rheumatoid arthritis
 - Active infectious disease such as HIV, HBV, active COVID, active (not treated) HCV
 - Obesity (BMI > 35) is a relative exclusion
 - BMI < 18.5
 - Any chemotherapy or radiation for any condition, including cancer treatment in any organ
(patients may be included with untreated or surgically only treated cancer in another organ)
 - Metastatic malignancy
 - Whole blood transfusion in the past 48 hours
 - Inflammatory bowel disease, microscopic colitis, and infectious colitis
 - Known inherited conditions with increased risk of colon cancer such as Lynch Syndrome, FAP, DNA repair defects etc
 - Use of blood thinners other than a baby aspirin (due to increased bleeding risk)
 - Underlying liver disease with cirrhosis or coagulopathy (due to increased bleeding risk)
- Note: while diverticulosis and colon polyps are not exclusions, we will not collect samples within 10 cm of these lesions.

Collection Protocol

1. U54 CRC will coordinate with GI team, to screen for colonoscopy cases where normal tissue may be available.
2. Scheduled in advance, U54 CRC or BRPC team member will approach patient's in-person for consent.

3. If patient/parents' consent, U54 CRC will coordinate with the GI Physician on procedure timing and biopsy procurement.
4. Upon day of procedure, Receive maximum 8 forceps biopsy fragments, 3-4 from the mid-ascending colon ("right colon") and 3-4 from the mid-descending colon (or the sigmoid colon as alternate site) ("left colon").
5. Process the biopsy fragments into the following samples. Keep samples from right colon and left colon separate.
 - a. FFPE – 1 biopsy per block
 - b. Snap frozen:
 - i. 1 snap freeze in cryovial as normal
 - ii. 1 will be put into freezing media
 1. Wash tissue with PBS to remove the blood and fecal matter.
 2. Place biopsy fragment into cryopreserved tubes with 1mL freezing media.
 3. Cells should be frozen slowly at 1°C/min. Put tubes in the freezing container (e.g., Mr. Frosty) with isopropyl alcohol and place them overnight in a -80°C freezer.
 4. Next day, transfer tubes to liquid nitrogen.
6. Route FFPE samples to Research Histology Lab to:
 - a. Create an H&E from each FFPE sample and route to pathologist for review.
 - b. After review, create 2 tissue scrolls from the tissue blocks and coordinate RNA and DNA extraction.
 - c. Once RNA and DNA extraction is complete, the frozen LN2 samples will be shipped to lab for scRNAseq