VERSION 2

AUG 08, 2019

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.56qg9dw

External link:
<https://mmpc.org/shared/document.aspx?id=256&docType=Protocol>

Protocol Citation: Jennifer Rutkowski 2019. UC Davis - Immunohistochemistry Factor VIII. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.56qg9dw>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: Aug 05, 2019

Last Modified: Aug 08, 2019

PROTOCOL integer ID:
 26544

UC Davis - Immunohistochemistry Factor VIII V.2

Jennifer Rutkowski¹

¹University of California, Davis

Mouse Metabolic Phenotyping Centers
 Tech. support email: info@mmpc.org

Lili Liang

ABSTRACT

Summary:

Factor VIII is a glycoprotein present in human plasma, human endothelial cells, megakaryocytes and platelets. Immunohistochemical staining for factor VIII related antigen could be used to determine if the benign and malignant neoplastic lesions are of endothelial origin. Furthermore, it can be utilized to determine vessel density in a tissue of interest

Modified from: IHC Methods and Materials VMTH - Anatomic Pathology, UC-Davis.

MATERIALS

MATERIALS

- ☒ xylene **Contributed by users**
- ☒ ethanol **Contributed by users**
- ☒ Hydrogen peroxide **Contributed by users**
- ☒ Methanol **Contributed by users**
- ☒ Proteinase K **Dako Catalog #S3020**
- ☒ 0.1M Phosphate Buffered Saline pH 7.4 **Contributed by users**
- ☒ Normal horse serum **Contributed by users**
- ☒ Tween-20 **Contributed by users**
- ☒ Factor VIII Ab **Dako Catalog #A0082**
- ☒ biotin-avidin based HRP **Biocare Medical Catalog #GR608**
- ☒ NovaRed for peroxidase **Vector Laboratories Catalog #SK-4800**
- ☒ Streptavidin-HRP HP604 **Biocare Medical Catalog #HP604**
- ☒ Mayer's Hematoxylin **Dako Catalog #S3309**

Keywords:

Immunohistochemistry Factor
VIII

coverslip **Corning Catalog #2935-245**

Reagent Preparation:**Reagent1: PBS-Tween 20**

Reagents and Materials:

0.1M Phosphate buffered saline (PBS), pH 7.4

Tween 20

Procedure:

For 100mL combine 100mL PBS and 20uL Tween 20

Reagent 2: Antibody diluent/blocking solution

Reagents and Materials:

PBS-Tween 20

Normal Horse Serum (NHS)

Procedure:

For 100mL combine 90mL PBS-Tween 20 and 10mL NHS.

Note:

Biocare Medical, [RRID: SCR-013549](#)

Vector Laboratories, [RRID:SCR_000821](#)

Dako [RRID:SCR_013530](#)

Factor VIII Ab # A0082, Cite this, (**Agilent Cat# A0082**, [RRID:AB_2315602](#))

SAFETY WARNINGS **WARNING:**

Formalin is, toxic, flammable and considered a carcinogen

Xylene, ethanol and methanol are all flammable and should be used in fume hood away from open flames or sparks

All blood components and biological materials should be handled as potentially hazardous. Follow universal precautions established by CDC when handling and disposing of infectious agents.

- 1 Immunohistochemistry was performed on four-micron thick, formalin-fixed, paraffin-embedded tissue sections, mounted on charged slides, and air-dried overnight at 37° C.
- 2 Sections were deparaffinized through xylene to 100% reagent alcohol, and then treated with 0.3% hydrogen peroxide in 100% methanol for 30 minutes.
- 3 Sections were rehydrated to deionized water through 95% and 70% reagent alcohols. Antigen retrieval was performed on sections for IBA-1 with heat induced epitope retrieval in a Black & Decker Steamer using Target Retrieval Solution, pH 6 (Dako S1699) for 30 minutes at 95°C, followed by a 20 minute cool down.
- 4 Retrieval was performed on sections for Factor VIII with Proteinase K (Dako S3020) at room temperature for 10 minutes.
- 5 After antigen retrieval, slides were rinsed in deionized water and placed in 0.1M Phosphate Buffered Saline, pH 7.4 (PBS). The antibody diluent and blocking reagent were PBS-Tween 20 (0.02%) and 10% normal horse serum (NHS) in PBS-Tween 20, respectively.
- 6 Sections were blocked for 20 minutes with antibody diluent and primary antibodies were applied without rinsing and incubated for 1 hour.
 - a. All post-antigen retrieval incubations are in a humidity chamber at room temperature.
- 7 After primary incubation, samples are rinsed twice for three minutes with PBS-Tween 20 between each subsequent reagent application.
- 8 A single step, polymer based HRP (BioCare Medical, RC542H) was applied for 30 minutes to label rabbit anti-IBA-1. A dual step, biotin-avidin based HRP (Biocare Medical, 4+ Detection System GR608) was applied for 10 minutes to link rabbit anti-Factor VIII.
- 9 Streptavidin-HRP (Biocare Medical HP604) was applied for 10 minutes to label the biotin link.

- 10 All labels were visualized with NovaRed for peroxidase (Vector SK-4800), per manufacturer's instructions.
- 11 Sections are counterstained in Mayer's Hematoxylin, air dried and coverslipped.