

May 15, 2019

## UC Davis - Lipoprotein profiling and partical size

DOI

[dx.doi.org/10.17504/protocols.io.yrifv4e](https://dx.doi.org/10.17504/protocols.io.yrifv4e)



Peter Havel<sup>1</sup>

<sup>1</sup>University of California, Davis

Mouse Metabolic Phenotyping Centers  
Tech. support email: [info@mmpc.org](mailto:info@mmpc.org)



Lili Liang

OPEN  ACCESS



DOI: [dx.doi.org/10.17504/protocols.io.yrifv4e](https://dx.doi.org/10.17504/protocols.io.yrifv4e)

External link: <https://mmpc.org/shared/document.aspx?id=263&docType=Protocol>

**Protocol Citation:** Peter Havel 2019. UC Davis - Lipoprotein profiling and partical size. **protocols.io**  
<https://dx.doi.org/10.17504/protocols.io.yrifv4e>

**License:** This is an open access protocol distributed under the terms of the **Creative Commons Attribution License**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

**Protocol status:** Working

**We use this protocol and it's working**

**Created:** March 01, 2019

**Last Modified:** May 15, 2019

**Protocol Integer ID:** 21002

**Keywords:** Lipoprotein profiling, Lipoprotein size

## Abstract

### Summary:

Lipoprotein profiling and Lipoprotein size will be provided for CM, VLDL, LDL and HDL as well as 20 additional subclasses of lipoprotein. Testing requires as little as 10 µL for full analytical testing.

### Before start

Lipoprotein profiling and size determined by LipoSEARCH <http://www.lipo-search.com/eng/index.php>

- 1 The sample is loaded into a gel permeation column specifically designed for separating lipoprotein components. Then, lipoprotein is eluted from the column in order of larger particles to smaller ones.
- 2 Lipoprotein fractionated by particle size is fed into reaction coils, and the degraded products after reacting with reagents are sent to detectors. The levels of cholesterol and triglyceride are output in the form of chromatogram.
- 3 The data obtained is processed with our patented analyzing program. The final output is a composite chromatogram and numeric data on 4 major classes (CM, VLDL, LDL and HDL) and 20 subclasses.