MAY 15, 2019

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.yrrfv56

External link:
<https://mmpc.org/shared/document.aspx?id=107&docType=Protocol>

Protocol Citation: John Rutledge 2019. UC Davis - Macrovascular Permeability and Lipoprotein Flux.
protocols.io
<https://dx.doi.org/10.17504/protocols.io.yrrfv56>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: Mar 01, 2019

Last Modified: May 15, 2019

PROTOCOL integer ID:
 21009

🌐 UC Davis - Macrovascular Permeability and Lipoprotein Flux

John Rutledge¹

¹University of California, Davis

Mouse Metabolic Phenotyping Centers

Tech. support email: info@mmpc.org

Lili Liang

ABSTRACT

Summary:

One of the three indices of arterial function that are compromised to a varying degree in individuals with cardiovascular disease is vascular permeability. This assay measures vascular permeability (as flux of labeled large molecular weight molecules: i.e. albumin or dextran) and lipid permeability (as flux of labeled lipid) in coronary or carotid arteries.

Modified from: Walsh et. al. Arterioscler Thromb Vasc Biol. 1999 Apr;19(4):840-6.

Keywords: cardiovascular disease, vascular permeability, lipid permeability

MATERIALS

MATERIALS

☒ Krebs-Henseleit Solution Contributed by users Catalog #See Below

☒ FITC- Dextran Sigma Aldrich Catalog #FD4, FD40S, or FD70

☒ TRITC- Dextran Sigma Aldrich Catalog #T1037 or T1162

☒ FITC- Albumin Sigma Aldrich Catalog #A9771

☒ TRITC- Albumin Sigma Aldrich Catalog #A2289

☒ Alexa-546 label Sigma Aldrich Catalog #10237

☒ DiI labeled Lipid Contributed by users Catalog #See protocol

☒ pentobarbital Cardinal Health

☒ DMEM Invitrogen - Thermo Fisher Catalog #11885

Reagent Preparation:

Reagent 1: Krebs-Henseleit Solution

116 mM NaCl, 5 mM KCl, 2.4 mM $\text{CaCl}_2 \cdot \text{H}_2\text{O}$, 1.2 mM MgCl_2 , 1.2 mM NH_2PO_4 , and 11mM glucose

Note:

Sigma-Aldrich, [RRID:SCR_008988](#)

SAFETY WARNINGS

WARNING:

All blood components and biological materials should be handled as potentially hazardous. Follow universal precautions established by CDC when handling and disposing of infectious agents.

- 1 Mice are anesthetized with an intraperitoneal injection with 50 mg pentobarbital/kg weight.
- 2 Blood was collected from each animal through the right atrium using a 22-gauge needle and a heparinized syringe. Blood was transferred to sterile Vacutainers and centrifuged at 2800 rpm for 10 min. Plasma samples were separated from blood cells and kept at 4°C.
- 3 Vasculature was flushed with DMEM by infusion into the left ventricle of heart.

- 4 Either common carotid arteries or aorta are dissected and cannulated.
 - a. An anterior midline skin incision was made from the mandible to the sternum and superficial neck muscles were retracted and the carotid arteries were carefully dissected free from surrounding tissue
 - b. An incision was made in the proximal artery and a cannula (polyethylene-50 tubing) was inserted and tied into place with 4-0 silk
 - c. A second incision was made in the distal portion of the artery, just proximal to the bifurcation of the common carotid artery, and another cannula was inserted and tied into place
- 5 Artery is perfused with Krebs's-Henseleit solution plus 1% bovine serum albumin gassed with 95% compressed air and 5% CO₂ until the start of the experiment.

Note: During our experiments, the artery is maintained at a constant length from the moment it is cannulated until the time it (via the cannulae) is secured into place in the perfusion chamber.
- 6 Cannulated artery was placed in a clear fluid-filled superfusate chamber and mounted on a Nikon MM-11 upright microscope stage for viewing.
- 7 Perfusate flowed through the artery at a physiological flow rate (1.5-2ml/min) and hydrostatic pressure (90 cm H₂O). All perfusates are maintained at 37°C and pH 7.3-7.4. Perfusate included:
 - a. Clear, non-fluorescent solution of Krebs's-Henseleit solution with 1% BSA
 - b. FITC- or TRITC- labeled Dextran or Albumin at 40-70 µg/mL in Krebs's-Henseleit solution with 1% BSA
 - c. DiI (see previous protocol) or Alex-546 (according to manufacturer's instructions) labeled lipid particles at 50 µg/mL in Krebs's-Henseleit solution with 1% BSA.
- 8 The artery in the superfusate chamber is brought into focus using a Nikon Plan X4 objective (NA 0.1). Fluorophore is excited and emission (see chart below) measured by a Nikon P1 photometer and input to a chart recorder and computer and visualized by Hamamatsu CCD television and camera.

Label	Excitation	Emission
DiL	549	565
FITC	495	521
TRITC	494	518
Alexa-546	554	570

- 9 The artery is perfused with the nonfluorescent solution to determine a baseline level of fluorescence intensity (If).

- 10 The artery is then perfused with the buffer solution containing the fluorescently labeled compound (FLC) for 10 min, immediately followed by wash out (10 min) of the fluorescent solution with the clear non-fluorescent solution.
 - a. Permeability: FITC-or TRITC- labeled albumin or dextran
 - b. Lipoprotein Flux: Dil or Alex546 labeled lipoproteins

- 11 If₀ represents the initial fluorescence intensity of the solution filling the artery lumen and If_A represents change in fluorescent intensity between baseline (If) and after washout (accumulation) all measures in millivolts (mV).

- 12 **Calculation of flux (ng/min/cm²)**

$$\text{Artery}_V = \pi r^2 l \quad \text{Artery}_{SA} = 2\pi r l$$

$$\text{Artery}_V * [\text{FLC in ng/mL}] = \text{ng in vessel at If}_0$$

$$\text{ng in vessel at If}_0 / \text{If}_0 \text{ (mV)} = \text{ng/mV}$$

$$\text{ng/mV} * \text{If}_A \text{ (mV)} = \text{ng compound at If}_A$$

$$\text{ng of compound at If}_A / \text{time(min)} / \text{Artery}_{SA} = \text{ng/min/cm}^2$$